

STRING TOPOLOGY, EULER CLASS AND TNCZ FREE LOOP FIBRATIONS.

LUC MENICHI

ABSTRACT. Let M be a connected, closed oriented manifold. Let $\omega \in H^m(M)$ be its orientation class. Let $\chi(M)$ be its Euler characteristic. Consider the free loop fibration $\Omega M \xrightarrow{i} LM \xrightarrow{ev} M$. For any class $a \in H^*(LM)$ of positive degree, we prove that the cup product $\chi(M)a \cup ev^*(\omega)$ is null. In particular, if $i^* : H^*(LM; \mathbb{F}_p) \rightarrow H^*(\Omega M; \mathbb{F}_p)$ is onto then $\chi(M)$ is divisible by p (or M is a point).

1. INTRODUCTION

There are few real applications of String Topology in algebraic topology: the author knows only one due to Kallel and Salvatore [14]. In this paper, we give another one.

Denote by $LM := \text{map}(S^1, M)$ the free loop space on M . Except where specified, we work over an arbitrary principal ideal domain \mathbb{k} .

In String Topology, shriek maps are used to defined operations. But usually shriek maps are used to obtained vanishing results: see for example [2, III.10.1] for an application of the transfer map in group cohomology. In this paper, after defining them carefully, we use the operations in String Topology to obtain the following vanishing result:

Theorem 1. *(Theorem 27 3) and Remark 32 below) Let M be a connected, closed oriented manifold. Let $\omega \in H^m(M)$ be its orientation class. Let $\chi(M)$ be its Euler characteristic. The cohomology of the free loops relative to the constant loops $H^*(LM, M)$ satisfies*

$$H^*(LM, M) \cup \chi(M)ev^*(\omega) = \{0\}.$$

This vanishing result also holds for any generalized cohomology h^* and homotopy fibre product of (the pull-back) of an embedding with itself (Theorem 12 4) below).

Using Leray-Hirsch theorem or Serre spectral sequence, we deduce

Corollary 2. *(Corollary 39 below) Let M be a connected, closed oriented manifold. Suppose that the free loop fibration $\Omega M \xrightarrow{i} LM \xrightarrow{ev} M$*

1991 *Mathematics Subject Classification.* 55P50, 55R40.

Key words and phrases. String Topology, free loop space, Euler Class.

is *Totally Non-Cohomologous to Zero* with respect to a field \mathbb{F} , i. e. $H^*(i; \mathbb{F}) : H^*(LM; \mathbb{F}) \twoheadrightarrow H^*(\Omega M; \mathbb{F})$ is onto. Then $\chi(M) = 0$ in \mathbb{F} or M is a point.

Again this Corollary is generalized for any generalized cohomology h^* and any fibration with section (Lemmas 15 and 19 below). We deduce then the following theorem.

Theorem 3. *(Theorem 21) Let $g : G \hookrightarrow E$ be the pull-back of an embedding in the sense of definition 8. Under some mild hypothesis, if the fibration p_g associated to g is *Totally Non-Cohomologous to Zero* and if all the homotopy fibres $p_g^{-1}(*)$ are not acyclic then the Euler class of g is null.*

In the case of the diagonal embedding, Theorem 3 gives Corollary 2.

We now give the plan of the paper.

Part 1. We construct carefully the shriek maps used in String Topology to define the operations. In particular, we give the key property (Proposition 6) that we use in this paper.

Part 2. We give our most general results. In section 5, we define the open string product and open string coproduct of Sullivan [26]. In section 6, we compute the open string coproduct of the homotopy fibre product of an embedding with itself. In section 7, we give general results on *Totally Non-Cohomologous to Zero* fibrations with sections using Leray-Hirsch (Lemma 15) or Serre spectral sequence (Lemma 19). These general results are used to prove Theorem 3. In section 8, as an example, we consider the case when the embedding is the inclusion of complex projective spaces.

Part 3. We specialize to the case of free loop spaces where the embedding is the diagonal embedding. In section 9, we define the Chas-Sullivan loop product [3] and the loop coproduct. In section 10, we compute the dual of the loop coproduct in term of cup product. In particular, we recover the results of Tamanoi [29] and Sullivan [26] concerning the vanishing of the loop coproduct. In section 11, we give (Theorem 33) a variant of Theorem 1. As application, we prove an homotopy version of a classical result relating fixed point action of the circle and Euler characteristic. And we prove Corollary 2. In section 12, we give many examples showing that Corollary 2 is pertinent. We conjecture that Corollary 2 holds for any simply-connected finite CW-complex. In section 13, we consider the case of relative free loop spaces. This is an example where the embedding is the pull-back of the diagonal embedding. As an application, we generalize Theorem 1 to the space $\text{map}(\vee_n S^1, M)$ of maps from the wedge of n circles to M .

Acknowledgment: We wish to thank our expert in generalized cohomology, Geoffrey powell. We would like also to thank the universities of Angers and Nantes for giving us the opportunity to teach a Master course on characteristic classes following [22].

Part 1. The shriek maps

2. THE SHRIEK MAP OF AN ORIENTED EMBEDDING

Let h^* be a generalized cohomology theory which is additive and multiplicative.

Let $\phi : M \hookrightarrow B$ an embedding between two manifolds without boundary of dimensions m and b respectively. Following [22, Corollary 11.2], we suppose that $\phi(M)$ is a closed subset of B , i. e. [19, Proposition A.53 (c), Theorem A.57] Φ is proper. Of course, this is the case if M is compact.

We also suppose that the normal bundle ν is h^* -oriented. If M and B are both h^* -oriented, ν is h^* -oriented since TM and $TB|_M = TM \oplus \nu$ are h^* -oriented ([12, Chapter 4, Lemma 4.1] or [7, Theorem 6 p. 45]).

By the tubular neighborhood theorem ([22, 11.1] or [12, Chapter 4, Theorem 5.2]), there exists an open neighborhood V of M in B and a diffeomorphism $exp : \nu \xrightarrow{\cong} V$ such that under this diffeomorphism, the zero section map $M \rightarrow \nu$ corresponds to the inclusion map $s : M \hookrightarrow V$.

Consider the associated closed disk bundle $D(\nu)$ and the associated sphere bundle $S(\nu)$. Let $N := exp(D(\nu))$ be a closed tubular neighborhood. Let $\partial N := exp(S(\nu))$ be its boundary. Note that the inclusion map $s : M \xrightarrow{\cong} N$ is a homotopy equivalence.

Since ν is h^* -oriented, there exists a Thom class $u \in h^{b-m}(D(\nu); S(\nu))$ and a Thom isomorphism [22, Theorem 9.1].

Remark 4. Our generalized cohomology h^* does not necessarily satisfies the weak equivalence axiom. Therefore the Thom homomorphism might not be an isomorphism [30, (17.9.1)].

The Thom class u will be thought as an element of $h^{b-m}(N; \partial N)$.

Since M is closed in B , $(B, B - M, N)$ is an excisive triad. The inclusion $\partial N \xrightarrow{\cong} N - M$ is a homotopy equivalence. Therefore the composite

$$i : (N, \partial N) \rightarrow (N, N - M) \rightarrow (B, B - M)$$

induces an isomorphism in cohomology. Let $j : B \rightarrow (B, B - M)$ be the canonical map. By definition [8, p. 419], $\phi^!$ is the composite

$$h^*(M) \xrightarrow[\cong]{s^{*-1}} h^*(N) \xrightarrow[\cong]{-\cup u} h^{*+b-m}(N, \partial N) \xrightarrow[\cong]{i^{*-1}} h^{*+b-m}(B, B - M) \xrightarrow{j^*} h^{*+b-m}(B)$$

3. THE SHRIEK MAP OF THE PULL-BACK OF AN EMBEDDING

The idea to construct the shriek map of the pull-back of an embedding, is to pull-back the Thom class and the tubular neighborhood. In particular, we will forget the original vector bundle ν and the fact the Thom homomorphism was (may-be see Remark 4) a Thom isomorphism.

Consider a (Serre) fibration $p : E \rightarrow B$. Consider the pull-back diagram

$$\begin{array}{ccc} \tilde{M} & \xrightarrow{\tilde{\phi}} & E \\ q \downarrow & & \downarrow p \\ M & \xrightarrow{\phi} & B \end{array}$$

The goal of this section is to construct a shriek map for $\tilde{\phi}$.

Let $\tilde{N} := p^{-1}(N)$. Then $p^{-1}(N - M) = p^{-1}(N) - p^{-1}(M) = \tilde{N} - \tilde{M}$. Let $\partial\tilde{N} := p^{-1}(\partial N)$. Consider the two rectangles where all the squares are pull-backs

$$\begin{array}{ccccc} \tilde{M} & \xrightarrow{\tilde{s}} & \tilde{N} & \longrightarrow & E \\ q \downarrow & & p \downarrow & & \downarrow p \\ M & \xrightarrow{s} & N & \longrightarrow & B \end{array} \quad \begin{array}{ccccc} \partial\tilde{N} & \xrightarrow{\tilde{}} & \tilde{N} - \tilde{M} & \longrightarrow & E \\ p \downarrow & & \downarrow & & \downarrow p \\ \partial N & \xrightarrow{\tilde{}} & N - M & \longrightarrow & B \end{array}$$

Since the inclusion map $s : M \xrightarrow{\sim} N$ is a homotopy equivalence and $p : \tilde{N} \rightarrow N$ is a (Serre) fibration, $\tilde{s} : \tilde{M} \xrightarrow{\sim} \tilde{N}$ is a (weak) homotopy equivalence. Similarly, since the inclusion $\partial N \xrightarrow{\sim} N - M$ is a homotopy equivalence, the inclusion $\partial\tilde{N} \xrightarrow{\sim} \tilde{N} - \tilde{M}$ is also a (weak) homotopy equivalence.

Since the inverse image of an excisive triad is an excisive triad, $(E, E - \tilde{M}, \tilde{N}) = (p^{-1}(B), p^{-1}(B - M), p^{-1}(N))$ is an excisive triad. Therefore the composite

$$\tilde{i} : (\tilde{N}, \partial\tilde{N}) \rightarrow (\tilde{N}, \tilde{N} - \tilde{M}) \rightarrow (E, E - \tilde{M})$$

induces an isomorphism in cohomology.

Let \tilde{u} be the image of the Thom class u by $p^* : h^*(N, \partial N) \rightarrow h^*(\tilde{N}, \partial\tilde{N})$. Let $\tilde{j} : E \rightarrow (E, E - M)$ be the canonical map. By definition, $\tilde{\phi}^!$ is the composite

$$h^*(\tilde{M}) \xrightarrow{\tilde{s}^{*-1}} h^*(\tilde{N}) \xrightarrow{-\cup\tilde{u}} h^{*+b-m}(\tilde{N}, \partial\tilde{N}) \xrightarrow{\tilde{i}^{*-1}} h^{*+b-m}(E, E - \tilde{M}) \xrightarrow{\tilde{j}^*} h^{*+b-m}(E)$$

Comparing with the definition of the shriek map of ϕ given in Section 2, since $\tilde{u} := p^*(u)$, we obviously have the naturality with respect

to pull-backs:

$$(5) \quad p^* \circ \phi^! = \tilde{\phi}^! \circ q^*.$$

Till now, our construction of the shriek map $\tilde{\phi}$ follows the construction of Tamanoi in the special case of the loop coproduct [28] and of the loop coproduct [29] in string topology, except that Tamanoi, in each case, construct a specific homotopy equivalence $\tilde{N} \xrightarrow{\sim} \tilde{M}$ replacing our homotopy equivalence $\tilde{s} : \tilde{M} \xrightarrow{\sim} \tilde{N}$. As remarked by Tamanoi [28, p. 8], note that in order to define $\tilde{\phi}^!$, we don't need to know if the total space $q^*(S(\nu))$ of the bundle induced by pulling-back $S(\nu)$, is diffeomorphic [25, Proposition 5.3], homeomorphic [6, p. 8], or homotopy equivalent to $\partial\tilde{N}$. So, we don't prove it in this note.

4. THE EULER CLASS

Proposition 6. *For the shriek map of the pull-back of an embedding ϕ , we have the formula for any $x \in h^*(\tilde{M})$:*

$$\tilde{\phi}^* \circ \tilde{\phi}^!(x) = x \cup q^*(e_\nu).$$

Here e_ν is the Euler class of the normal bundle ν of the embedding ϕ .

For an embedding $\phi : M \hookrightarrow B$, this formula is well known [15, Theorem 6.1 (5)]. For $\tilde{\phi}$, the pull-back of an embedding, the proof will be similar [15, p. 282].

Proof. (See [27, Theorem 2.1 (5)] for a different proof) Let $\tilde{s}_{rel} : \tilde{M} \rightarrow (\tilde{N}, \partial\tilde{N})$ be the relative inclusion map. Remark that the following square commutes

$$\begin{array}{ccc} (\tilde{N}, \partial\tilde{N}) & \xrightarrow{\tilde{i}} & (E, E - \tilde{M}) \\ \tilde{s}_{rel} \uparrow & & \uparrow \tilde{j} \\ \tilde{M} & \xrightarrow{\tilde{\phi}} & E. \end{array}$$

Remark also that

$$\tilde{s}_{rel}^* : h^*(\tilde{N}, \partial\tilde{N}) \rightarrow h^*(\tilde{M})$$

is $h^*(\tilde{N})$ -linear where $h^*(\tilde{N})$ acts on $h^*(\tilde{M})$ by restriction of scalar with respect to the algebra morphism $\tilde{s}^* : h^*(\tilde{N}) \rightarrow h^*(\tilde{M})$. Therefore, by

definition of $\tilde{\phi}^!$,

$$\begin{aligned}\tilde{\phi}^* \circ \tilde{\phi}^!(x) &= \tilde{\phi}^* \circ \tilde{j}^* \circ \tilde{i}^{*-1}(\tilde{s}^{*-1}(x) \cup \tilde{u}) \\ &= \tilde{s}_{rel}^* \circ \tilde{i}^* \circ \tilde{i}^{*-1}(\tilde{s}^{*-1}(x) \cup \tilde{u}) \\ &= (\tilde{s}^* \circ \tilde{s}^{*-1}(x)) \cup \tilde{s}_{rel}^*(\tilde{u}) = x \cup \tilde{s}_{rel}^*(\tilde{u})\end{aligned}$$

Consider the commutative diagram

$$\begin{array}{ccc}\tilde{M} & \xrightarrow{\tilde{s}_{rel}} & (\tilde{N}, \partial\tilde{N}) \\ q \downarrow & & \downarrow p \\ M & \xrightarrow{s_{rel}} & (N, \partial N)\end{array}$$

Since, by definition, $\tilde{u} := p^*(u)$, we obviously have

$$\tilde{s}_{rel}^*(\tilde{u}) = q^* \circ s_{rel}^*(u).$$

Note that, by definition ([15, p. 279] or [22, p. 98]), $s_{rel}^*(u)$ is the Euler class e_ν of the normal bundle ν . \square

By naturality of the Euler class [22, Property 9.2], note that $q^*(e_\nu)$ is the Euler class of the vector bundle $q^*(\nu)$ induced by pulling-back ν along $q : \tilde{M} \rightarrow M$.

Corollary 7. *Let $k : F \hookrightarrow E$ be the inclusion of the fiber of the fibration p . If $b > m$ then $k^* \circ \tilde{\phi}^! = 0$ in singular cohomology.*

Proof. Let $l : F \hookrightarrow \tilde{M}$ be the inclusion of the fiber of the induced fibration q . We have the commutative diagram

$$\begin{array}{ccccc} & & k & & \\ & & \curvearrowright & & \\ F & \xrightarrow{l} & \tilde{M} & \xrightarrow{\tilde{\phi}} & E \\ \varepsilon \downarrow & & q \downarrow & & \downarrow p \\ \{*\} & \xrightarrow{\eta} & M & \xrightarrow{\phi} & B\end{array}$$

where the two squares are pull-backs. By Proposition 6,

$$\tilde{\phi}^* \circ \tilde{\phi}^!(x) = x \cup q^*(e_\nu)$$

where $e_\nu \in H^{b-m}(M)$ is the Euler class of the normal bundle ν . Therefore

$$k^* \circ \tilde{\phi}^!(x) = l^* \circ \tilde{\phi}^* \circ \tilde{\phi}^!(x) = l^*(x) \cup l^* \circ q^*(e_\nu) = l^*(x) \cup \varepsilon^* \circ \eta^*(e_\nu) = 0.$$

\square

Part 2. The general case

5. THE OPEN STRING (CO)PRODUCTS

In this section, we define the joining product \wedge of Sullivan [26] that following Tamanoi [27], we prefer to call the open string product. And more important for us, we define a generalized version of the cutting at time 1/2 product $\vee_{1/2}$ of Sullivan [26] that following Tamanoi [27], we prefer to call the open string coproduct.

Let $f : F \rightarrow E$ and $g : G \rightarrow E$ be two maps. Let

$${}^f E^g = \{(a, \omega, b) \in F \times E^I \times G / f(a) = \omega(0), g(b) = \omega(1)\}$$

denote the *homotopy fibre product* of f and g which is obtained by the following pull-back

$$\begin{array}{ccc} {}^f E^g & \longrightarrow & E^I \\ (ev_0, ev_1) \downarrow & & \downarrow (ev_0, ev_1) \\ F \times G & \xrightarrow{(f \times g)} & E \times E. \end{array}$$

Definition 8. A continuous map $g : G \rightarrow E$ is the *pull-back of an embedding* if it is equipped with a pull-back diagram

$$\begin{array}{ccc} G & \xrightarrow{g} & E \\ q \downarrow & & \downarrow p \\ M & \xrightarrow{\phi} & B \end{array}$$

where $\phi : M \hookrightarrow B$ a proper embedding between two manifolds of codimension m with h^* -oriented normal bundle and where the map p is a (Serre) fibration (This is exactly the case considered in Section 3).

Remark 9. Suppose that a continuous map $g : G \rightarrow E$ fits into a pull-back diagram

$$\begin{array}{ccc} G & \xrightarrow{g} & E \\ q \downarrow & & \downarrow p \\ M & \xrightarrow{\phi} & B \end{array}$$

where $\phi : M \hookrightarrow B$ a proper embedding between two manifolds of codimension m with h^* -oriented normal bundle and where the map p is smooth and is transverse to ϕ .

Then $g : G \hookrightarrow E$ a proper embedding between two manifolds of codimension m with h^* -oriented normal bundle.

A common example to both definition 8 and remark 9 is when p is a smooth fibre bundle.

Proof of Remark 9. Since ϕ and p are transverse, $g : G \hookrightarrow E$ is an embedding of codimension m . Note that the normal bundle of g , ν_g , is the pull-back $q^*(\nu_\phi)$ of the normal bundle of ϕ , ν_ϕ , along q . Therefore since ν_ϕ is h^* -oriented, ν_g is also h^* -oriented [1, 11.7.11] without assuming that G or E is h^* -orientable. Since $\phi(M)$ is a closed subset of B , $g(G) = p^{-1}(\phi(M))$ is a closed subset of E . \square

Theorem 10. [26] *Let $f : F \rightarrow E$, $g : G \rightarrow E$ and $h : H \rightarrow E$ be three maps.*

1) *If G is a smooth h^* -oriented manifold without boundary of dimension n then there is a open string product*

$$lp_{f,g,h} : H_*({}^f E^g) \otimes H_*({}^g E^h) \rightarrow H_{*-n}({}^f E^h).$$

2) *Suppose that g is the pull-back of an embedding in the sense of definition 8. Then there is a open string coproduct*

$$lcp_{f,g,h} : h^*({}^f E^g) \otimes h^*({}^g E^h) \rightarrow h^{*+m}({}^f E^h).$$

Remark 11. In [26], Sullivan consider the open string product only in the case when g is an embedding.

Proof. 1) Consider the three pull-back squares (Compare with [27, Diagram p.11])

$$\begin{array}{ccccc} {}^f E^g \times {}^g E^h & \xleftarrow{\tilde{\Delta}} & {}^f E^g \times_G {}^g E^h & \xrightarrow{\mu_{f,g,h}} & {}^f E^h \\ \text{ev}_1 \times \text{ev}_0 \downarrow & & \text{ev}_{1/2} \downarrow & & \downarrow \text{ev}_{1/2} \\ G \times G & \xleftarrow{\Delta} & G & \xrightarrow{g} & E \\ & & q \downarrow & & p \downarrow \\ & & M & \xrightarrow{\phi} & B \end{array}$$

The open string product $lp_{f,g,h}$ is defined as the Chas-Sullivan loop product above using the left pull-back square:

$$H_* {}^f E^g \otimes H_* {}^g E^h \xrightarrow{\times} H_* {}^f E^g \times {}^g E^h \xrightarrow{\tilde{\Delta}_!} H_{*-n} {}^f E^g \times_G {}^g E^h \xrightarrow{\mu_{f,g,h}^*} H_{*-n} {}^f E^h.$$

2) Since p is a (Serre) fibration, then the composite

$${}^f E^h \xrightarrow{\text{ev}_{1/2}} E \xrightarrow{p} B$$

is also a fibration. Therefore using the total right rectangle, since $\phi : M \hookrightarrow B$ is an embedding, we obtain the shriek map

$$\mu_{f,g,h}^! : h^*({}^f E^g \times_G {}^g E^h) \rightarrow h^{*+m}({}^f E^h).$$

The open string coproduct $lcp_{f,g,h}$ is now defined as the composite

$$h^*({}^f E^g) \otimes h^*({}^g E^h) \xrightarrow{\times} h^*({}^f E^g \times {}^g E^h) \xrightarrow{\tilde{\Delta}^*} h^*({}^f E^g \times_G {}^g E^h) \xrightarrow{\mu_{f,g,h}^1} h^{*+m}({}^f E^h).$$

□

6. A SIMPLE FORMULA FOR THE OPEN STRING COPRODUCT

Theorem 12. *Let $f : F \rightarrow E$, $g : G \rightarrow E$ and $h : H \rightarrow E$ be three maps. Suppose that g is the pull-back of an embedding in the sense of definition 8. Let $e_\nu \in h^m(M)$ be the Euler class of the normal bundle of the embedding $\phi : M \hookrightarrow B$.*

Let $ev_0 : {}^g E^h \rightarrow G$, $(a, w, b) \mapsto a$ be the first projection map. Let $ev_1 : {}^f E^g \rightarrow G$, $(a, w, b) \mapsto b$ be the second projection map.

Let $\sigma : G \hookrightarrow {}^g E^g$, $b \mapsto (b, \text{constant path } g(b), b)$ be the section of both projection map ev_0 and ev_1 when $f = g = h$.

Then

1) *the open string coproduct*

$$lcp_{f,g,g} : h^*({}^f E^g) \otimes h^*({}^g E^g) \rightarrow h^{*+m}({}^f E^g)$$

is given by

$$lcp_{f,g,g}(a \otimes b) = a \cup ev_1^*(\sigma^*(b) \cup q^*(e_\nu)).$$

2) *the open string coproduct*

$$lcp_{g,g,h} : h^*({}^g E^g) \otimes h^*({}^g E^h) \rightarrow h^{*+m}({}^g E^h)$$

is given by

$$lcp_{g,g,h}(b \otimes c) = ev_0^* \circ \sigma^*(b) \cup c \cup ev_0^* \circ q^*(e_\nu).$$

3) *In the case $f = g = h$,*

$$ev_0^* \circ q^*(e_\nu) = ev_1^* \circ q^*(e_\nu) \in h^*({}^g E^g).$$

4) *The ideal $\text{Ker } \sigma^* : h^*({}^g E^g) \rightarrow h^*(G)$ satisfies*

$$\text{Ker } \sigma^* \cup ev_0^* \circ q^*(e_\nu) = \{0\}.$$

5) *the open string coproduct on $h^*({}^g E^g)$*

$$lcp_{g,g,g} : h^*({}^g E^g) \otimes h^*({}^g E^g) \rightarrow h^{*+m}({}^g E^g)$$

is given by

$$lcp_{g,g,g}(a \otimes b) = ev_0^*(\sigma^*(a) \cup \sigma^*(b) \cup q^*(e_\nu)) = ev_1^*(\sigma^*(a) \cup \sigma^*(b) \cup q^*(e_\nu)).$$

6) *Let $\alpha \in {}^g E^g$. Denote by ${}^g E_{[\alpha]}^g$ the path-connected component of α . Denote by ${}^g E_0^g := \cup_{\alpha \in G} {}^g E_{[\sigma(\alpha)]}^g$, i. e. the subspace of ${}^g E^g$, union of the path-connected components $\text{Im } \pi_0(\sigma) : \pi_0(G) \hookrightarrow \pi_0({}^g E^g)$ (If G is path-connected, there is only one). Then*

the open string coproduct $\text{lcp}_{f,g,g}$ is trivial outside of $h^*({}^f E^g) \otimes h^*({}^g E_0^g)$,
the open string coproduct $\text{lcp}_{g,g,h}$ is trivial outside of $h^*({}^g E_0^g) \otimes h^*({}^g E^h)$,
the open string coproduct $\text{lcp}_{g,g,g}$ is trivial outside of $h^*({}^g E_0^g) \otimes h^*({}^g E_0^g)$.

7) The open string coproduct $\text{lcp}_{f,g,g}$ maps $h^*({}^f E_{[\alpha]}^g) \otimes h^*({}^g E^g)$ to $h^{*+m}({}^f E_{[\alpha]}^g)$.

The open string coproduct $\text{lcp}_{g,g,h}$ maps $h^*({}^g E^g) \otimes h^*({}^g E_{[\alpha]}^h)$ to $h^{*+m}({}^g E_{[\alpha]}^h)$.

The image of the open string coproduct $\text{lcp}_{g,g,g}$ is contained in ${}^g E_0^g$.

8) Suppose that there is an integer $\dim G$ such that $\forall i > \dim G$, $h^i(G) = \{0\}$. Then for all $a, b \in h^*({}^g E^g)$ such that $|a| + |b| > \dim G - m$, $\text{lcp}_{g,g,g}(a \otimes b) = 0$.

Proof. 1) and 2) Recall that we have the following two pull-back squares

$$\begin{array}{ccc} {}^f E^g \times_G {}^g E^h & \xrightarrow{\mu_{f,g,h}} & {}^f E^h \\ \text{ev}_{1/2} \downarrow & & \downarrow \text{ev}_{1/2} \\ G & \xrightarrow{g} & E \\ q \downarrow & & \downarrow p \\ M & \xrightarrow{\phi} & B \end{array}$$

By Proposition 6, for any $x \in h^*({}^f E^g \times_G {}^g E^h)$

$$(13) \quad \mu_{f,g,h}^* \circ \mu_{f,g,h}^\dagger(x) = x \cup \text{ev}_{1/2}^* \circ q^*(e_\nu).$$

Note that till here, the same discussion for the open string product, shows that

$$(14) \quad \tilde{\Delta}^* \circ \tilde{\Delta}^\dagger(x) = x \cup \text{ev}_{1/2}^*(e_{TG}).$$

But now, we will see that both $\mu_{f,g,g}^*$ and $\mu_{g,g,h}^*$ admit a retract. In particular, $\mu_{g,g,g}^*$ admits two different retracts i_1^* and i_2^* . Let $i_1 : {}^f E^g \rightarrow {}^f E^g \times_G {}^g E^g$ be the inclusion map on the first factor defined by

$$i_1(a, w, b) = ((a, w, b), (b, \text{constant path } g(b), b))$$

for $a \in F$, $w \in E^I$ and $b \in G$ such that $f(a) = w(0)$ and $g(b) = w(1)$. Let $i_2 : {}^g E^h \rightarrow {}^g E^g \times_G {}^g E^h$ be the inclusion map on the second factor defined by

$$i_2(a, w, b) = ((a, \text{constant path } g(a), a), (a, w, b))$$

for $a \in G$, $w \in E^I$ and $b \in H$ such that $(g(a), h(b)) = (w(0), w(1))$. The first inclusion i_1 is a section up to homotopy of $\mu_{f,g,g}$. Therefore i_1^* is a retract of $\mu_{f,g,g}^*$.

STRING TOPOLOGY, EULER CLASS AND TNCZ FREE LOOP FIBRATIONS

So by formula (13), since $ev_{1/2} \circ i_1$ is the projection map $ev_1 : {}^f E^g \rightarrow G$, $(a, w, b) \mapsto b$, for any $x \in h^*({}^f E^g \times_G {}^g E^g)$,

$$\begin{aligned} \mu_{f,g,g}^!(x) &= i_1^* \circ \mu_{f,g,g}^* \circ \mu_{f,g,g}^!(x) = i_1^*(x \cup ev_{1/2}^* \circ q^*(e_\nu)) \\ &= i_1^*(x) \cup i_1^* \circ ev_{1/2}^* \circ q^*(e_\nu) = i_1^*(x) \cup ev_1^* \circ q^*(e_\nu). \end{aligned}$$

Similarly since i_2^* is a retract of $\mu_{g,g,h}^*$ and since $ev_{1/2} \circ i_2$ is the projection map $ev_0 : {}^g E^h \rightarrow G$, $(a, w, b) \mapsto a$, for any $x \in h^*({}^g E^g \times_G {}^g E^h)$,

$$\mu_{g,g,h}^!(x) = i_2^*(x) \cup ev_0^* \circ q^*(e_\nu).$$

Since the following two squares commute

$$\begin{array}{ccc} {}^f E^g & \xrightarrow{i_1} & {}^f E^g \times_G {}^g E^g & & {}^g E^h & \xrightarrow{i_2} & {}^g E^g \times_G {}^g E^h \\ \Delta \downarrow & & \downarrow \tilde{\Delta} & & \Delta \downarrow & & \downarrow \tilde{\Delta} \\ {}^f E^g \times {}^f E^g & \xrightarrow{id \times (\sigma \circ ev_1)} & {}^f E^g \times {}^g E^g & & {}^g E^h \times {}^g E^h & \xrightarrow{(\sigma \circ ev_0) \times id} & {}^g E^g \times {}^g E^h, \end{array}$$

for any $a \in h^*({}^f E^g)$ and $b \in h^*({}^g E^g)$,

$$\begin{aligned} \mu_{f,g,g}^! \circ \tilde{\Delta}^*(a \times b) &= i_1^* \circ \tilde{\Delta}^*(a \times b) \cup ev_1^* \circ q^*(e_\nu) \\ &= a \cup (\sigma \circ ev_1)^*(b) \cup ev_1^* \circ q^*(e_\nu) \end{aligned}$$

and for any $b \in h^*({}^g E^g)$ and $c \in h^*({}^g E^h)$,

$$\mu_{g,g,h}^! \circ \tilde{\Delta}^*(b \times c) = (\sigma \circ ev_0)^*(b) \cup c \cup ev_0^* \circ q^*(e_\nu).$$

3) Using 1),

$$l\text{cop}_{g,g,g}(1 \otimes 1) = ev_1^* \circ q^*(e_\nu).$$

Using 2),

$$l\text{cop}_{g,g,g}(1 \otimes 1) = ev_0^* \circ q^*(e_\nu).$$

4) Let ε be 0 or 1. Since $\sigma^* \circ ev_\varepsilon^* = id$, using the split short exact sequence

$$0 \rightarrow \text{Ker } \sigma^* \xrightarrow{i} h^*({}^g E^g) \xrightarrow{\sigma^*} h^*(G) \rightarrow 0,$$

we obtain that $\text{Ker } \sigma^* = \text{Im } r_\varepsilon$ where $r_\varepsilon : h^*({}^g E^g) \rightarrow \text{Ker } \sigma^*$ is the retract of the inclusion defined by $r_\varepsilon(a) = a - ev_\varepsilon^* \circ \sigma^*(a)$.

By 1) and 3),

$$l\text{cop}_{g,g,g}(a \otimes 1) = a \cup ev_1^* \circ q^*(e_\nu) = a \cup ev_0^* \circ q^*(e_\nu).$$

By 2),

$$l\text{cop}_{g,g,g}(a \otimes 1) = ev_0^* \circ \sigma^*(a) \cup ev_0^* \circ q^*(e_\nu).$$

Therefore $r_0(a) \cup ev_0^* \circ q^*(e_\nu) = 0$.

5) By 4),

$$b \cup ev_0^* \circ q^*(e_\nu) = ev_0^* \circ \sigma^*(b) \cup ev_0^* \circ q^*(e_\nu).$$

Therefore by 2)

$$lcop_{g,g,g}(a \otimes b) = ev_0^* \circ \sigma^*(a) \cup ev_0^* \circ \sigma^*(b) \cup ev_0^* \circ q^*(e_\nu).$$

By 4) and 3)

$$a \cup ev_1^* \circ q^*(e_\nu) = ev_1^* \circ \sigma^*(a) \cup ev_1^* \circ q^*(e_\nu).$$

Therefore by 1) and the graded commutativity of the cup product

$$lcop_{g,g,g}(a \otimes b) = ev_1^* \circ \sigma^*(a) \cup ev_1^* \circ \sigma^*(b) \cup ev_1^* \circ q^*(e_\nu).$$

6) Since by definition, σ arrives inside ${}^g E_0^g$, σ^* factorizes through the projection $h^*({}^g E^g) \rightarrow h^*({}^g E_0^g)$. So using 1) and 2), 6) is proved.

7) Let $a \in h^*({}^f E_{[\alpha]}^g)$ and $b \in h^*({}^g E^g)$. The cohomology of a space is isomorphic to the product of the cohomology algebras of its path-connected components. Therefore the cup product with $ev_1^*(\sigma^*(b) \cup q^*(e_\nu))$ defines a linear application from $h^*({}^f E_{[\alpha]}^g)$ to itself. So by 1), $lcop_{f,g,g}(a \otimes b) \in h^*({}^f E_{[\alpha]}^g)$.

In the case $f = g$, by 6), if $lcop_{g,g,g}(a \otimes b)$ is non-zero then the path-connected component of α belongs to the image of $\pi_0(\sigma)$ and in this case $lcop_{g,g,g}(a \otimes b) \in h^*({}^f E_0^g)$.

8) The element $\sigma^*(a) \cup \sigma^*(b) \cup q^*(e_\nu) \in h^*(G)$ is of degree $|a| + |b| + m > \dim G$ and so is null. Using 5), we obtain that $lcop_{g,g,g}(a \otimes b) = 0$. \square

7. TNCZ FIBRATIONS

Lemma 15. *Suppose that B is path-connected. Suppose also that B is a CW-complex or that h^* satisfies the weak equivalence axiom. Let $F \xrightarrow{i} E \xrightarrow{p} B$ be a (Serre) fibration with base B which admits a section $\sigma : B \rightarrow E$ up to homotopy, i. e. $p \circ \sigma \approx id_B$ and an element $e \in h^*(B)$ such that*

$$Ker \sigma^* \cup p^*(e) = \{0\}.$$

If the fibration p is Totally Non-Cohomologous to Zero, i. e. $h^(i) : h^*(E) \rightarrow h^*(F)$ is onto and if $\tilde{h}^*(F)$ is a finitely generated free graded h^* -module then $e = 0$ or $\tilde{h}^*(F) = \{0\}$.*

Remark 16. Let $p : E \rightarrow B$ be a fibration. Let $\omega : I \rightarrow B$ be a path from b to b' . Let $w_\# : p^{-1}(b) \xrightarrow{\cong} p^{-1}(b')$ be the induced homotopy equivalence between the fibers [24, Theorem 2.8.12]. By definition, $w_\#$ commutes up to homotopy with the inclusions of fibers $i_b : p^{-1}(b) \hookrightarrow E$ and $i_{b'} : p^{-1}(b') \hookrightarrow E$. In cohomology, $i_b^* = w_\#^* \circ i_{b'}^*$ [30, Proof of (17.9.3)]. So i_b^* is surjective if and only if $i_{b'}^*$ is surjective. And given a family of vectors $c_j \in h^*(E)$, the $i_b^*(c_j)$'s form a h^* -basis of $h^*(p^{-1}(b))$ if and only if the $i_{b'}^*(c_j)$'s form a h^* -basis of $h^*(p^{-1}(b'))$.

Proof. Let $b_0 \in B$. Let $\varepsilon_F : F \rightarrow *$ be the unique map to a point. Since B is path-connected and $\tilde{h}^*(F) := \text{coker } \varepsilon_F^* : h^* \rightarrow h^*(F)$ does not depend of a base point and is homotopy invariant [30, 17.1.3], we can chose any fibre F . We take $F := p^{-1}(\{p \circ \sigma(b_0)\})$. Since $h^*(i) : h^*(E) \rightarrow h^*(F)$ is surjective, $\tilde{h}^*(i) : \tilde{h}^*(E) \rightarrow \tilde{h}^*(F)$ is also surjective.

Since $\tilde{h}^*(\sigma) \circ \tilde{h}^*(p) = id$, $\tilde{h}^*(E) = \text{Ker } \tilde{h}^*(\sigma) \oplus \text{Im } \tilde{h}^*(p)$. Since $\text{Im } \tilde{h}^*(p) \subset \text{Ker } \tilde{h}^*(i)$, the restriction of $\tilde{h}^*(i)$ to $\text{Ker } \tilde{h}^*(\sigma)$ is also surjective.

Suppose that $\tilde{h}^*(F) \neq \{0\}$. Then there exists classes $c_j \in \text{Ker } \tilde{h}^*(\sigma) \subset \tilde{h}^*(E)$ such that the $\tilde{h}^*(i)(c_j)$'s form a h^* -basis of $\tilde{h}^*(F)$.

Now let $b_0 \in B$, $\sigma(b_0) \in E$ and $\sigma(b_0) \in F$ be the chosen base points $*$ of B , E and F . Denote by $\eta_{x_0}^X : \{x_0\} \hookrightarrow X$ be the inclusion of the base point x_0 into a based space X . Then we have the canonical identification for a based space X between $\tilde{h}^*(X)$ and $h^*(X, x_0) \cong \text{Ker } \eta_{x_0}^{X*} : h^*(X) \rightarrow h^*$. Since $\sigma : (B, *) \hookrightarrow (E, *)$ and $i : (F, *) \hookrightarrow (E, *)$ are based maps, we can consider that the classes $c_j \in h^*(E, *)$, that the $i^*(c_j)$'s form a h^* -basis of $h^*(F, *)$, that $\text{Ker } \tilde{h}^*(\sigma) = \text{Ker } \sigma^*$ and so that $c_j \cup p^*(e) = 0$.

Since $h^*(F) \cong h^*(F, *) \oplus h^*$, the classes $c_j \in h^*(E, *)$ and the unit $1 \in h^0(E)$ are send by i^* to a h^* -basis of $h^*(F)$. So by the Leray-Hirsch theorem for generalized cohomology ([30, (17.8.4)] using Remark 16), $h^*(E)$ is a free $h^*(B)$ -modules with basis 1 and the c_j 's. So $e = 0$. \square

Remark 17. In Lemma 15, when the generalized cohomology h^* is a singular cohomology H^* , it is enough to suppose that $\tilde{H}^q(F)$ is \mathbb{k} -free module of finite type for each degree $q \geq 0$. Indeed in this case, we can apply the Leray-Hirsch theorem for singular cohomology [11, Exercise 3 p. 51] (see also [30, (17.8.1)] using Remark 16 or [23, Theorem 4.4] where the fibre of the fibration is assumed to be path-connected). In Lemma 19 below, we improve further Lemma 15 for singular cohomology. The proof of Lemma 19 relies on the following interesting Lemma:

Lemma 18. *Let $F \xrightarrow{i} E \xrightarrow{p} B$ be a (Serre) fibration with path-connected base B which admits a section $\sigma : B \rightarrow E$ up to homotopy, i. e. $p \circ \sigma \approx id_B$ and an element $e \in H^m(B)$ such that*

$$\text{Ker } \sigma^* \cup p^*(e) = \{0\}.$$

Consider the cohomological Serre spectral sequence $(E_r^{,*}, d_r)$ associated to the fibration p . Suppose that the action of $\pi_1(B)$ on $H^*(F)$ is trivial.*

Let f be an element of $\tilde{H}^*(F)$. Suppose that f is in the image of $i^* : H^*(E) \rightarrow H^*(F)$.

Denote by $e \otimes 1$, the image of e by the canonical morphism [23, III.2.10.10]

$$i_* : H^m(B) \rightarrow H^m(B; H^0(F)) = E_2^{m,0}$$

Denote by $1 \otimes f$ the image of f by the inverse of the canonical morphism [23, III.2.10.9]

$$i_0^* : E_2^{0,*} = H^0(B; H^*(F)) \xrightarrow{\cong} H^0(*; H^*(F)) = H^*(F).$$

Then the product $e \otimes 1 \cup 1 \otimes f \in E_2^{m,*}$ must be killed: there exist $r \geq 2$ and $x \in E_r^{*,*}$ such that $d_r(x) = e \otimes 1 \cup 1 \otimes f$.

Proof. Since B is path-connected, the triviality of the local coefficients $H^*(F)$ implies that i_0^* is an isomorphism (and conversely by [23, III.1.18 (3)]).

For degree reasons, $\forall r \geq 2, d_r(e \otimes 1) = 0$ and $e \otimes 1$ in $E_\infty^{m,0} = F^m H^m(E)$ is $p^*(e)$. Since $f \in \text{Im } i^*, \forall r \geq 2, d_r(1 \otimes f) = 0$. Let q be the degree of f . Let $c \in \tilde{H}^q(E)$ such that $\tilde{H}^*(i)(c) = f$. As explained in the proof of Lemma 15, c can be chosen in $\text{Ker } \sigma^*$. Then $1 \otimes f$ in $E_\infty^{0,q} = H^q(E)/F^1 H^q(E)$ is the class of c . Therefore $e \otimes 1 \cup 1 \otimes f$ in $E_\infty^{m,q} = F^m H^{m+q}(E)/F^{m+1} H^{m+q}(E)$ is the class of $p^*(e) \cup c$. Since $\text{Ker } \sigma^* \cup p^*(e) = \{0\}, p^*(e) \cup c = 0$. Therefore $e \otimes 1 \cup 1 \otimes f$ must be killed. \square

Lemma 19. Let $F \xrightarrow{i} E \xrightarrow{p} B$ be a (Serre) fibration with path-connected base B which admits a section $\sigma : B \rightarrow E$ up to homotopy, i. e. $p \circ \sigma \approx \text{id}_B$ and an element $e \in H^m(B)$ such that

$$\text{Ker } \sigma^* \cup p^*(e) = \{0\}.$$

Suppose that $\forall n \in \mathbb{N}, H_n(B)$ is a finitely generated \mathbb{k} -module or $\forall q \in \mathbb{N}, H^q(F)$ is a finitely generated \mathbb{k} -module. Suppose also that $\forall q \geq 2, H^q(F)$ is a torsion free \mathbb{k} -module.

If the fibration p is Totally Non-Cohomologous to Zero, i. e. $H^*(i) : H^*(E) \rightarrow H^*(F)$ is onto and if $H^m(B)$ is a free \mathbb{k} -module then $e = 0$ or $\tilde{H}^*(F) = \{0\}$.

Proof. Since $H^*(i)$ is onto, the action of $\pi_1(B)$ on $H^*(F)$ is trivial [23, III.Theorem 4.4].

By hypothesis, $H^q(F)$ is a finitely generated \mathbb{k} -module or $\forall n \geq 0, H_n(B)$ is a finitely generated \mathbb{k} -module. So since \mathbb{k} is a principal ideal domain by [24, Theorem 5.5.10], we have a short exact sequence

$$0 \rightarrow H^p(B) \otimes H^q(F) \xrightarrow{\mu} E_2^{p,q} \rightarrow \text{Tor}^{\mathbb{k}}(H^{p+1}(B), H^q(F)) \rightarrow 0$$

where μ is a morphism of algebras. Therefore, since $\forall q \geq 0$ $H^q(F)$ is torsion free, $\forall p, q \in \mathbb{N}$ $E_2^{p,q} \cong H^p(B) \otimes H^q(F)$ as algebras. Since i^* is onto, d_r is null on $E^{0,q}$. Therefore the Serre spectral sequence collapses on the E_2 -term (Here we have reproved the well-known [23, III.Theorem 4.4] with weaker hypothesis).

So by Lemma 18, for any $f \in \tilde{H}^q(F)$, the element $e \otimes f \in E^{m,q} = H^m(B) \otimes H^q(F)$ must be zero. Since $H^m(F)$ is free, $e \otimes f = 0$ implies that $e = 0$ or f has torsion. \square

The following lemma is a generalization of Lemma 19 if the base B of the fibration is not path-connected.

Lemma 20. *Let $B = \cup_{\beta \in \pi_0(B)} B_\beta$ the decomposition of B into its path-connected components. Let $p : E \rightarrow B$ be a (Serre) fibration with base B which admits a section $\sigma : B \rightarrow E$ up to homotopy, i. e. $p \circ \sigma \approx id_B$ and an element $e = (e_\beta)_{\beta \in \pi_0(B)} \in H^m(B) = \prod_{\beta \in \pi_0(B)} H^m(B_\beta)$ such that*

$$\text{Ker } \sigma^* \cup p^*(e) = \{0\}.$$

Denote by F_β , the fibre $p^{-1}(b)$ when $b \in B_\beta$. For all $\beta \in \pi_0(B)$, suppose that $\forall n \in \mathbb{N}$, $H_n(B_\beta)$ is a finitely generated \mathbb{k} -module or $\forall q \in \mathbb{N}$, $H^q(F_\beta)$ is a finitely generated \mathbb{k} -module. Suppose also that $\forall \beta \in \pi_0(B)$, $\forall q \geq 2$, $H^q(F_\beta)$ is a torsion free \mathbb{k} -module.

If the fibration p is Totally Non-Cohomologous to Zero, i. e. $\forall \beta \in \pi_0(B)$ $H^(i_\beta) : H^*(E) \rightarrow H^*(F_\beta)$ is onto and if $\forall \beta \in \pi_0(B)$ $H^m(B_\beta)$ is a free \mathbb{k} -module then $\forall \beta \in \pi_0(B)$ ($e_\beta = 0$ or $\tilde{H}^*(F_\beta) = \{0\}$).*

Proof. For all $\beta \in \pi_0(B)$, we apply Lemma 19 to the fibration

$$F_\beta \xrightarrow{i_\beta} E_{\pi_0(p)^{-1}(\beta)} \xrightarrow{p_\beta} B_\beta$$

obtained by restricting p to the union $E_{\pi_0(p)^{-1}(\beta)}$ of path-connected components α of E such that $\pi_0(p)(\alpha) = \beta$. The fibration p_β admits the restriction of σ to B_β , $\sigma_\beta : B_\beta \rightarrow E_{\pi_0(p)^{-1}(\beta)}$ as section up to homotopy. The product of maps

$$\prod_{\beta \in \pi_0(B)} \sigma_\beta^* : \prod_{\beta \in \pi_0(B)} H^*(E_{\pi_0(p)^{-1}(\beta)}) \rightarrow \prod_{\beta \in \pi_0(B)} H^*(B_\beta)$$

can be identified with $\sigma^* : H^*(E) \rightarrow H^*(B)$. Therefore $\text{Ker } \sigma^* \cup p^*(e)$ can be identified with $\prod_{\beta \in \pi_0(B)} (\text{Ker } \sigma_\beta^* \cup p_\beta^*(e_\beta))$. \square

Of course, the following theorem generalizes the fundamental Corollary 11.4 of [22].

Theorem 21. *Let $g : G \hookrightarrow E$ be the pull-back of an embedding in the sense of definition 8. Let $e_\nu \in h^m(M)$ be the Euler class of the normal bundle of the embedding $\phi : M \hookrightarrow B$. Let $p_g : E^I \times_g G \rightarrow E$ be the fibration associated to g defined by $p_g((\omega, b)) = \omega(0)$ for any $b \in G$ and*

any path $\omega : I \rightarrow E$ such that $\omega(1) = g(b)$. Let $p_\phi : B^I \times_\phi M \rightarrow B$ be the fibration associated to ϕ .

Suppose that $\forall n \in \mathbb{N}$, $H_n(G)$ is a finitely generated \mathbb{k} -module or $\forall y \in B$, $\forall q \in \mathbb{N}$, $H^q(p_\phi^{-1}(y))$ is a finitely generated \mathbb{k} -module. Suppose also that $\forall y \in B$, $\forall q \geq 2$, $H^q(p_\phi^{-1}(y))$ is a torsion free \mathbb{k} -module.

If the fibration p_g is Totally Non-Cohomologous to Zero, i. e. $\forall x \in E$ $H^*(i_x) : H^*(E^I \times_g G) \rightarrow H^*(p_g^{-1}(x))$ is onto and if $H^m(G)$ is a free \mathbb{k} -module then $\forall b \in G$ either $\tilde{H}^*(p_\phi^{-1}(p \circ g(b))) = \{0\}$ or the component of $q^*(e_\nu)$ in $H^m(G_{[b]})$ the cohomology of the path-connected component of b in G is trivial.

Proof. By definition of the homotopy fibre product ${}^f E^g$, for any $a \in F$, we have the following commutative diagram of spaces

$$\begin{array}{ccccc}
 ev_0^{-1}(a) & \xrightarrow{\cong} & p_g^{-1}(f(a)) & \xrightarrow{\cong} & p_\phi^{-1}(p \circ f(a)) \\
 \downarrow & & \downarrow i_{f(a)} & & \downarrow \\
 {}^f E^g & \longrightarrow & E^I \times_g G & \xrightarrow{p^I \times q} & B^I \times_\phi M \xleftarrow[\approx]{s} M \\
 ev_0 \downarrow & & p_g \downarrow & & p_\phi \downarrow \\
 F & \xrightarrow{f} & E & \xrightarrow{p} & B
 \end{array}$$

$\swarrow \phi$

where the bottom left square is a pull-back, s is a homotopy equivalence and the bottom right square is a homotopy pull-back (i. e. the induced map $p_g^{-1}(f(a)) \rightarrow p_\phi^{-1}(p \circ f(a))$ between the homotopy fibre of g and ϕ is a homotopy equivalence). Since $\forall a \in F$, $H^*(i_{f(a)}) : H^*(E^I \times_g G) \rightarrow H^*(p_g^{-1}(f(a)))$ is onto, $H^*({}^f E^g) \rightarrow H^*(ev_0^{-1}(a))$ is also onto, i. e. ev_0 is Totally Non-Cohomologous to Zero.

Suppose now that $f = g$ (and $F = G$). By part 4) of Theorem 12,

$$\text{Ker } \sigma^* \cup ev_0^* \circ q^*(e_\nu) = \{0\}.$$

By applying Lemma 20 to the fibration $ev_0 : {}^g E^g \rightarrow G$ we obtain that for all $b \in G$, the component of $q^*(e_\nu)$ in $H^*(G_{[b]})$ is trivial or $\tilde{H}^*(ev_0^{-1}(b)) = \{0\}$. \square

8. AN EXAMPLE

Corollary 22. *Consider the following pull-back diagram*

$$\begin{array}{ccc}
 G & \xrightarrow{g} & E \\
 q \downarrow & & \downarrow p \\
 \mathbb{C}P^q & \xrightarrow{\phi} & \mathbb{C}P^n
 \end{array}$$

where $p : E \rightarrow \mathbb{C}\mathbb{P}^n$ is a (Serre) fibration over the n -th complex projective space $\mathbb{C}\mathbb{P}^n$ and $\phi : \mathbb{C}\mathbb{P}^q \hookrightarrow \mathbb{C}\mathbb{P}^n$ is the inclusion, $0 \leq q < n$.

If the fibration p_g associated to g is Totally Non-Cohomologous to Zero and if $H^{2n-2q}(G)$ is a free \mathbb{k} -module then $q^*(a^{n-q}) = 0$. Here a is a generator of $H^2(\mathbb{C}\mathbb{P}^q)$.

Proof. By [22, Theorem 14.10], $c(T\mathbb{C}\mathbb{P}^n)$, the total Chern class of the tangent bundle of $\mathbb{C}\mathbb{P}^n$ is equal to $(1+a)^{n+1}$ in $H^\Pi(\mathbb{C}\mathbb{P}^n)$. Since $T\mathbb{C}\mathbb{P}^q \oplus \nu = T\mathbb{C}\mathbb{P}^n|_{\mathbb{C}\mathbb{P}^q}$, in $H^\Pi(\mathbb{C}\mathbb{P}^q)$,

$$c(\nu) = c(T\mathbb{C}\mathbb{P}^n|_{\mathbb{C}\mathbb{P}^q})/c(T\mathbb{C}\mathbb{P}^q) = (1+a)^{n+1}/(1+a)^{q+1} = (1+a)^{n-q}.$$

Therefore $e(\nu) = c_{n-q}(\nu) = a^{n-q}$.

We have a morphism of S^1 -principal fibre bundles

$$\begin{array}{ccc} S^{2q+1} & \xrightarrow{\tilde{\phi}} & S^{2n+1} \\ \downarrow & & \downarrow \\ \mathbb{C}\mathbb{P}^q & \xrightarrow{\phi} & \mathbb{C}\mathbb{P}^n \end{array}$$

where $\tilde{\phi} : S^{2q+1} \hookrightarrow S^{2n+1}$ is the inclusion. Since this square is a pull-back, the homotopy fibre of ϕ , $p_\phi^{-1}(*)$, is homotopy equivalent to the homotopy fibre of $\tilde{\phi}$. Since $\pi_{2q+1}(S^{2n+1}) = \{0\}$, $\tilde{\phi}$ is homotopically trivial and its homotopy fibre is homotopy equivalent to $S^{2q+1} \times \Omega S^{2n+1}$. Therefore by Theorem 21, all the components of $q^*(e_\nu)$ are trivial. \square

Corollary 23. *Let $f : \mathbb{C}\mathbb{P}^p \hookrightarrow \mathbb{C}\mathbb{P}^n$ and $g : \mathbb{C}\mathbb{P}^q \hookrightarrow \mathbb{C}\mathbb{P}^n$ be the inclusions, $0 \leq p < n$, $0 \leq q < n$. If the fibration $ev_0 : {}^f E^g \rightarrow \mathbb{C}\mathbb{P}^p$ is Totally Non-Cohomologous to Zero then $q \geq p$ and $n > p + q$.*

Proof. Suppose that $q < p$. Since $ev_0^* \circ f^* = ev_1^* \circ g^*$ and f^* is surjective, for all $q < i \leq p$,

$$ev_0^*(a^i) = ev_0^* \circ f^*(a^i) = ev_1^* \circ g^*(a^i) = ev_1^*(a^i) = 0.$$

Therefore ev_0^* is not injective and so by [23, III.Theorem 4.4], ev_0 is not Totally Non-Cohomologous to Zero.

Consider the Serre spectral sequence associated to the fibration $F \hookrightarrow {}^f E^g \xrightarrow{ev_1} \mathbb{C}\mathbb{P}^q$. We saw in the proof of Corollary 22 that its fibre F is homotopy equivalent to $S^{2p+1} \times \Omega S^{2n+1}$. Therefore $H^+(F)$ is concentrated in degree $\geq 2p + 1$. And so $E_r^{s,t} \neq \{0\} \Rightarrow t = 0$ or $t \geq 2p + 1$. Therefore ev_1^* is an isomorphism in degree $\leq 2p$. (In particular, if $p \geq q$, ev_1^* is injective).

Suppose now that ev_0 is Totally Non-Cohomologous to Zero and that $n - q \leq p$. Then $H^{2n-2q}(ev_1) : H^{2n-2q}(\mathbb{C}\mathbb{P}^q) \xrightarrow{\cong} H^{2n-2q}({}^f E^g)$ is an isomorphism. Since $H^{2n-2q}(\mathbb{C}\mathbb{P}^q)$ is \mathbb{k} -free, by Corollary 22,

$H^{2n-2q}(ev_1)(a^{n-q}) = 0$. And so $a^{n-q} = 0$ in $H^*(\mathbb{C}\mathbb{P}^q)$. Therefore $n - q > q$. In particular, $p > q$. \square

Remark 24. In the case $p = q$ of Corollary 23, parts 4) and 8) of Theorem 12 give that

$$\text{Ker } \sigma^* \cup ev_0^*(a^{n-q}) = \{0\}$$

and that for all $b \in H^*({}^g E^g)$ of degree $> 4q - 2n$, $b \cup ev_0^*(a^{n-q}) = 0$.

Remark 25. (Over \mathbb{Q} , the converse of Corollary 23 is true) Over \mathbb{Q} , a relative Sullivan model of ev_0 is given by the inclusion of differential graded algebras

$$(\Lambda(x_2, z_{2p+1}), d) \hookrightarrow (\Lambda(x_2, z_{2p+1}, t_{2q+1}, sy_{2n+1}), d)$$

with $d(z_{2p+1}) = x_2^{p+1}$, $d(t_{2q+1}) = x_2^{q+1}$ and $d(sy_{2n+1}) = t_{2q+1}x_2^{n-q} - z_{2p+1}x_2^{n-p}$ (Compare with [21, Example 7.3]).

If $n > p + q$, by replacing sy by $sy - ztx^{n-p-q-1}$, we can assume that $d(sy) = 0$. If $q \geq p$, by replacing t by $t - zx^{q-p}$, we can assume that $d(t) = 0$. Therefore if $n > p + q$ and $q \geq p$ then over \mathbb{Q} , ev_0 is Totally Non-Cohomologous to Zero.

Part 3. the free loops case

In this part, we consider our main example of homotopy fibre product, the space LM of free loops on a manifold.

9. THE LOOP PRODUCT AND THE LOOP COPRODUCT

Let M be a smooth oriented manifold without boundary. In this section, M is not necessarily compact. The diagonal map $\Delta : M \hookrightarrow M \times M$ is an embedding. Since M is Hausdorff, $\Delta(M)$ is a closed subset of $M \times M$. As we have explained in Section 2, we can define the shriek map of Δ , $\Delta_!$ in homology.

By definition, the *intersection product* in homology, is the composite

$$H_*(M) \otimes H_*(M) \xrightarrow{\times} H_*(M \times M) \xrightarrow{\Delta_!} H_{*-m}(M).$$

We have the following push-out squares

$$\begin{array}{ccccc} S^1 & \amalg & S^1 & \longrightarrow & S^1 \vee S^1 & \xleftarrow{c} & S^1 \\ & \uparrow & & & \uparrow & & \uparrow \\ \star & \amalg & \star & \longrightarrow & \star & \longleftarrow & S^0 \end{array}$$

where $c : S^1 \rightarrow S^1 \vee S^1$ is the comultiplication or pinch map of S^1 . Note that all the vertical maps are cofibrations. Since the functor

$map(-, M)$ transforms push-out squares in pull-back squares, we have the following pull-back squares where all the vertical maps are fibrations

$$\begin{array}{ccccc}
 LM \times LM & \xleftarrow{\tilde{\Delta}} & LM \times_M LM & \xrightarrow{\mu} & LM \\
 ev \times ev \downarrow & & q \downarrow & & \downarrow (ev, ev_{1/2}) \\
 M \times M & \xleftarrow{\Delta} & M & \xrightarrow{\Delta} & M \times M
 \end{array}$$

and $\mu := map(c, M)$ is the composition or multiplication of loops. Since $\Delta : M \hookrightarrow M \times M$ is an embedding, as we have explained in Section 3, we can define the shriek map of $\tilde{\Delta}$, $\tilde{\Delta}_!$ in homology, and the shriek maps of μ , $\mu_!$ in homology, $\mu^!$ in cohomology.

By definition, the Chas-Sullivan loop product in homology, is the composite

$$H_*(LM) \otimes H_*(LM) \xrightarrow{\times} H_*(LM \times LM) \xrightarrow{\tilde{\Delta}_!} H_{*-m}(LM \times_M LM) \xrightarrow{\mu_*} H_{*-m}(LM).$$

By definition, the loop coproduct in homology is the composite

$$H_*(LM) \xrightarrow{\mu_*} H_{*-m}(LM \times_M LM) \xrightarrow{\tilde{\Delta}_*} H_{*-m}(LM \times LM).$$

In this note, we work over an arbitrary principal ideal domain \mathbb{k} and so the cross product is not in general an isomorphism. Therefore, we will consider the loop coproduct in cohomology. By definition, the loop coproduct in cohomology is the product defined by the composite

$$H^*(LM) \otimes H^*(LM) \xrightarrow{\times} H^*(LM \times LM) \xrightarrow{\tilde{\Delta}^*} H^*(LM \times_M LM) \xrightarrow{\mu^!} H^{*+m}(LM)$$

Remark 26. Let $k : \Omega M \hookrightarrow LM$ be the inclusion of the pointed loops into the free loops. If the dimension of M is positive, from Corollary 7, we obtain that the composite

$$H_*(\Omega M) \otimes H_*(\Omega M) \xrightarrow{k_* \otimes k_*} H_*(LM) \otimes H_*(LM) \xrightarrow{\text{loop product}} H_{*-m}(LM)$$

is trivial.

10. A SIMPLE FORMULA FOR THE LOOP COPRODUCT

Denote by $LM_{[1]}$ the path-connected component of LM of freely contractile loops. Recall that $ev : LM \rightarrow M$ is the evaluation map. Let $\sigma : M \hookrightarrow LM$, $m \mapsto$ constant loop m , be its trivial section.

Theorem 27. *Let M be a connected, closed \mathbb{k} -oriented manifold of dimension m . Let $\omega \in H^m(M)$ be its orientation class. Let $\chi(M)$ be its Euler characteristic. Then*

1) The loop coproduct, $\mu^! \circ \tilde{\Delta}^*$ on $H^*(LM)$ is given for $a, b \in H^*(LM)$, by

$$\mu^! \circ \tilde{\Delta}^*(a \otimes b) = \chi(M)a \cup ev^*(\sigma^*(b) \cup \omega).$$

Here \cup is the cup product on $H^*(LM)$.

2) The loop coproduct, $\mu^! \circ \tilde{\Delta}^*$ on $H^*(LM)$ is graded commutative with respect to the usual degrees: that is, for $a \in H^p(LM)$, $b \in H^q(LM)$

$$\mu^! \circ \tilde{\Delta}^*(a \otimes b) = (-1)^{pq} \mu^! \circ \tilde{\Delta}^*(b \otimes a).$$

3) The ideal $\text{Ker } \sigma^* : H^*(LM) \rightarrow H^*(M)$ satisfies

$$\text{Ker } \sigma^* \cup \chi(M)ev^*(\omega) = \{0\}.$$

4) The loop coproduct, $\mu^! \circ \tilde{\Delta}^*$ on $H^*(LM)$ is given for $a, b \in H^*(LM)$, by

$$\mu^! \circ \tilde{\Delta}^*(a \otimes b) = \chi(M)ev^*(\sigma^*(a) \cup \sigma^*(b) \cup \omega).$$

5) the loop coproduct, $\mu^! \circ \tilde{\Delta}^*$ on $H^*(LM)$ is trivial outside of $H^0(LM_{[1]}) \otimes H^0(LM_{[1]}) \cong \mathbb{k} \otimes \mathbb{k}$.

6) On $H^0(LM_{[1]}) \otimes H^0(LM_{[1]})$, the loop coproduct is given by

$$\mu^! \circ \tilde{\Delta}^*(1 \otimes 1) = \chi(M)ev^*(\omega).$$

7) The image of the loop coproduct $\mu^! \circ \tilde{\Delta}^*$ is contained in $H^*(LM_{[1]})$.

Remark 28. Over a field, parts 2), 5) and 7) of this Theorem are not new. Indeed over a field, the commutativity of the loop coproduct was proved by Cohen and Godin [5] and parts 5) and 7) are the duals of [29, Theorem B (2)].

Lemma 29. Consider ${}^\Delta M \times M^\Delta$ the self homotopy fibre product along the diagonal. Explicitly ${}^\Delta M \times M^\Delta$ is just the subspace

$$\{(\omega, \omega') \in M^I \times M^I / \omega(0) = \omega'(0), \omega(1) = \omega'(1)\}.$$

Let $\Theta : {}^\Delta M \times M^\Delta \xrightarrow{\cong} LM$ be the homeomorphism mapping (ω, ω') to the free loop $\omega * \omega'^{-1}$ obtained by composing the path ω with the inverse of the path ω' . Then

1) [26, Example iii) free loop space] With respect to the loop product and the open string product,

$$H_*(\Theta) : H_*({}^\Delta M \times M^\Delta) \xrightarrow{\cong} H_*(LM)$$

is an isomorphism of algebras.

2) With respect to the loop coproduct and the open string coproduct,

$$H^*(\Theta) : H^*(LM) \xrightarrow{\cong} H^*({}^\Delta M \times M^\Delta)$$

is an isomorphism of algebras.

STRING TOPOLOGY, EULER CLASS AND TNCZ FREE LOOP FIBRATIONS

Proof. Denote by $\rho_\alpha : LM \xrightarrow{\cong} LM$ the homeomorphism mapping a free loop l to the rotated free loop $t \mapsto l(t + \alpha)$. Up to the homeomorphism Θ , the two pull-back squares defining the open string (co)product on $\Delta M \times M^\Delta$

$$\begin{array}{ccccc} \Delta M \times M^\Delta \times \Delta M \times M^\Delta & \xleftarrow{\tilde{\Delta}} & \Delta M \times M^\Delta \times_{M \times M} \Delta M \times M^\Delta & \xrightarrow{\mu_{\Delta, \Delta, \Delta}} & \Delta M \times M^\Delta \\ \downarrow ev_1 \times ev_0 & & \downarrow ev_{1/2} & & \downarrow ev_{1/2} \\ M \times M & \xleftarrow{\Delta} & M & \xrightarrow{\Delta} & M \times M \end{array}$$

coincide with the following two vertical rectangles defining the loop (co)product since ρ_α is homotopic to the identity map.

$$\begin{array}{ccccc} LM \times LM & \longleftarrow & LM_{1/2} \times_M LM_0 & \longrightarrow & LM \\ \downarrow \rho_{1/2} \times id \cong & & \downarrow \cong & & \downarrow \rho_{1/4} \\ LM \times LM & \xleftarrow{\tilde{\Delta}} & LM \times_M LM & \xrightarrow{\mu} & LM \\ \downarrow ev \times ev & & \downarrow q & & \downarrow (ev, ev_{1/2}) \\ M \times M & \xleftarrow{\Delta} & M & \xrightarrow{\Delta} & M \times M \end{array}$$

□

Proof of Theorem 27. We apply Theorem 12 in the case where $f = g = h = \phi$ is the diagonal embedding $\Delta : M \hookrightarrow M \times M$.

The normal bundle ν of Δ is isomorphic to the tangent bundle of M , TM [22, Lemma 11.5]. Since M is compact and connected, the Euler class of the tangent bundle is the fundamental class multiplied by the Euler characteristic [22, Corollary 11.12]:

$$e_\nu = e_{TM} = \chi(M)\omega.$$

Using part 2) of Lemma 29, we have proved Theorem 27. □

Remark 30. Let M be a connected, non-compact \mathbb{k} -oriented manifold of dimension m and suppose that \mathbb{k} is a field. Then its loop coproduct $\mu^! \circ \tilde{\Delta}^*$ on $H^*(LM)$ is trivial.

Proof of Remark 30. Since M is non-compact then $H_m(M) = 0$. Since \mathbb{k} is a field, $H^m(M) = \text{Hom}_{\mathbb{k}}(H_m(M), \mathbb{k}) = 0$. So e_{TM} is trivial. Therefore the same proof as the proof of Theorem 27 shows that the loop coproduct is trivial.

Alternatively, for any $x \in H^*(LM \times_M LM)$

$$\mu^* \circ \mu^!(x) = x \cup q^*(e_{TM}) = 0.$$

Since the composition of loops μ admits a section, μ^* is injective and so $\mu^!$ is null. \square

Corollary 31. *The loop coproduct is trivial if and only if $\chi(M) = 0$ in \mathbb{k} .*

This corollary follows also from [29, (3-1) and (3-2)] (Compare also with [29, Corollary 3.2] or [26, Bottom p. 7]). In [4], Chataur and Thomas gave the first example of manifold with non-trivial loop coproduct.

Proof. Since $ev \circ \sigma = id$, ev^* is injective. Therefore since w is a basis of $H^m(M)$,

$$\chi(M)ev^*(w) = 0 \iff \chi(M)\omega = 0 \iff \chi(M) = 0 \text{ in } \mathbb{k}.$$

\square

Remark 32. Since $ev \circ \sigma = id_M$, M is a subspace of LM and we can consider the relative cohomology $H^*(LM, M)$. Using the long exact sequence associated, $H^*(LM, M)$ can be identified with $\text{Ker } \sigma^* : H^*(LM) \rightarrow H^*(M)$. From part 3) of Theorem 27, we have that the loop coproduct vanishes on $H^*(LM, M)$. In [26], Sullivan introduced a non trivial product on $H^*(LM, M)$ of different degree that he called the cutting at any time \vee . In [9], Goresky and Hingston rediscover this non trivial product that they denote \otimes .

11. APPLICATIONS

Theorem 33. *Let M be a connected, closed \mathbb{k} -oriented manifold of dimension m . Let $\omega \in H^m(M)$ be its orientation class. Let $\chi(M)$ be its Euler characteristic. Then*

- 1) $\chi(M)ev^*(\omega) \in H^m(LM_{[1]})$.
- 2) For any $a \in H^*(LM)$ of positive degree,

$$\chi(M)a \cup ev^*(\omega) = 0.$$

Proof. Comparing 6) and 7) in Theorem 27, we get 1).

By 5) and 1) in Theorem 27,

$$0 = \mu^! \circ \tilde{\Delta}^*(a \otimes 1) = \chi(M)a \cup ev^*(\omega).$$

\square

If \mathbb{k} is a field then 1) means that for all non contractile free loop α and for all $a \in H_m(LM_{[\alpha]})$,

$$\chi(M)H_m(ev)(a) = 0.$$

Remark 34. In general, $ev^*(\omega)$ does not belong to $H^*(LM_{[1]})$ and $a \cup ev^*(\omega)$ is not trivial: Suppose that \mathbb{k} is a field. Let G be a connected compact Lie group. Note that $\chi(G) = 0$. For any $[\alpha] \in \pi_1(G)$, let Θ_α be the usual isomorphism from the tensor product $H_*(\Omega_{[\alpha]}G) \otimes H_*(G)$ to $H_*(LG_{[\alpha]})$. Here $\Omega_{[\alpha]}G$ denotes the pointed loops of G homotopic to α . Let ε be the augmentation of $H_*(\Omega_{[\alpha]}G)$. The previous isomorphism Θ_α fits into the commutative triangle of graded vector spaces

$$\begin{array}{ccc} H_*(\Omega_{[\alpha]}G) \otimes H_*(G) & \xrightarrow[\cong]{\Theta_\alpha} & H_*(LG_{[\alpha]}) \\ & \searrow^{\varepsilon \otimes Id} \quad \swarrow_{H_*(ev)} & \\ & \mathbb{k} \otimes H_*(G) & \end{array}$$

Let $[G]$ be the fundamental class of G . Recall that $[\alpha]$ is a generator of $H_0(\Omega_{[\alpha]}G)$. Then

$$H_{\dim G}(ev) \circ \Theta_\alpha([\alpha] \otimes [G]) = [G] \neq 0$$

Therefore $ev^*(\omega)$ does not belong to $H^{\dim G}(LG_{[1]})$ for any non simply-connected, connected compact Lie group G (e. g. S^1).

Let $\eta : \mathbb{k} \rightarrow H^*(\Omega G)$ be the unit map of $H^*(\Omega G)$. The usual isomorphism of algebras Θ from the tensor product of graded algebras $H^*(\Omega G) \otimes H^*(G)$ to $H^*(LG)$ fits similarly into the commutative triangle of graded algebras

$$\begin{array}{ccc} H^*(\Omega G) \otimes H^*(G) & \xrightarrow[\cong]{\Theta} & H^*(LG) \\ & \searrow^{\eta \otimes Id} \quad \swarrow_{H^*(ev)} & \\ & \mathbb{k} \otimes H^*(G) & \end{array}$$

Therefore for any non-zero element b of $H^*(\Omega G)$,

$$\Theta(b \otimes 1) \cup H^*(ev)(\omega) = \Theta(b \otimes w) \neq 0.$$

If G is a connected compact Lie group such that $H^*(\Omega G)$ is not concentrated in degree 0 (e. g. S^3), we have obtained an element a of positive degree such that $a \cup ev^*(\omega)$ is non zero.

Corollary 35. *Let M be a connected, closed \mathbb{k} -oriented manifold of dimension m such that in \mathbb{k} , $\chi(M) \neq 0$. Let $\mu : S^1 \times M \rightarrow M$ be a continuous map such that the composite*

$$\{1\} \times M \rightarrow S^1 \times M \xrightarrow{\mu} M$$

is homotopic to the identity map. Then there exists an map $\nu : S^1 \times M \rightarrow M$ homotopic to μ who has at least a fixed point m_0 , i. e. $\nu(S^1 \times \{m_0\}) = m_0$.

Proof. Let $\sigma_\mu : M \rightarrow LM$ be the map sending $m \in M$ to its orbit $\mu(-, m) : S^1 \rightarrow M$. Since for all $m \in M$, $ev \circ \sigma_\mu(m) = \mu(1, m)$, σ_μ is a section up to homotopy of ev . Therefore in cohomology,

$$\sigma_\mu^* \circ ev^*(\chi(M)\omega) = \chi(M)\omega.$$

Since M is path-connected, σ_μ arrives in the path-connected component $LM_{[\alpha]}$ of a free loop α . So σ_μ^* is trivial outside of $H^*(LM_{[\alpha]})$. By 1) of Theorem 33, $\chi(M)ev^*(\omega) \in H^m(LM_{[1]})$. Since $\chi(M)$ is not zero in \mathbb{k} , $\chi(M)\omega = \sigma_\mu^* \circ ev^*(\chi(M)\omega)$ is not trivial. Therefore α is contractile, i. e. $[\alpha] = [1]$.

Let $i : \{m_0\} \hookrightarrow M$ be the inclusion of a non-degenerated base point into M . Since α is contractile, $\sigma_\mu(m_0)$ is homotopic to the constant loop \hat{m}_0 and so the following triangle commutes up to homotopy.

$$\begin{array}{ccc} S^1 \times M & \xrightarrow{\mu} & M \\ S^1 \times i \uparrow & \nearrow \hat{m}_0 & \\ S^1 \times \{m_0\} & & \end{array}$$

By the homotopy extension property of the cofibration $S^1 \times i : S^1 \times \{m_0\} \hookrightarrow S^1 \times M$, we can change up to homotopy μ into a map ν such that the triangle commutes now exactly. \square

Corollary 35 should be considered as an homotopy version of the following classic result:

Theorem 36. ([24, Theorem 4.7.12]. *Compare also with [15, Theorem 5.39 or Corollary 6.17]*) *Let M be a compact Euclidean Neighborhood Retract (e. g. a compact topological manifold [10, A.9]) such that $\chi(M) \neq 0$. Let $\mu : S^1 \times M \rightarrow M$ be an action of the circle on M . Then M has at least a fixed point.*

Remark 37. If a map $\mu : S^1 \times M \rightarrow M$ is only an action up to homotopy then it may happen that M has no fixed point. Therefore the conclusion of Corollary 35 cannot be improved in general: Consider the sphere $M = S^2$ in \mathbb{R}^3 . Let $\nu : S^1 \times S^2 \rightarrow S^2$ be the action given by rotation of axis z . Let $f : S^2 \rightarrow S^2$ be the rotation of angle π and of axis y . Since f is homotopic to the identity map, the composite $f \circ \nu$ is an action up to homotopy without any fixed point.

Corollary 38. *Let M be a connected, closed \mathbb{k} -oriented manifold of dimension m . Consider the cohomological Serre spectral sequence $(E_r^{*,*}, d_r)$ associated to the free loop fibration $\Omega M \xrightarrow{i} LM \xrightarrow{ev} M$. Suppose that the (conjugation) action of $\pi_1(M)$ on $H^*(\Omega M)$ is trivial.*

Let f be an element of $\tilde{H}^*(\Omega M)$. Suppose that f is in the image of $i^* : H^*(LM) \rightarrow H^*(\Omega M)$. Then $\chi(M)\omega \otimes f \in H^m(M) \otimes H^*(\Omega M) = E_2^{m,*}$ must be killed: there exist $r \geq 2$ and $x \in E_r^{*,*}$ such that $d_r(x) = \chi(M)\omega \otimes f$.

Proof. Since M is compact, $\forall n \geq 0$, $H_n(M)$ is a finitely generated \mathbb{k} -module. So since \mathbb{k} is a principal ideal domain by [24, Theorem 5.5.10], we have a short exact sequence

$$0 \rightarrow H^p(M) \otimes H^q(\Omega M) \xrightarrow{\mu} E_2^{p,q} \rightarrow \text{Tor}^{\mathbb{k}}(H^{p+1}(M), H^q(\Omega M)) \rightarrow 0$$

where μ is a morphism of algebras. Since $H^0(\Omega M)$, $H^1(M) \cong \text{Hom}(H_1(M), \mathbb{k})$ and $H^{m+1}(M) = 0$ are torsion free, $E_2^{p,q} \cong H^p(M) \otimes H^q(\Omega M)$ if $p = 0$ or $p = m$ or $q = 0$.

By part 3) of Theorem 27, $\text{Ker } \sigma^* \cup \chi(M)ev^*(\omega) = \{0\}$. So by Lemma 18, $\chi(M)\omega \otimes 1 \cup 1 \otimes f = \chi(M)\omega \otimes f$ must be killed. \square

Corollary 39. *Let M be a connected, closed \mathbb{k} -oriented manifold. Suppose that the free loop fibration $\Omega M \xrightarrow{i} LM \xrightarrow{ev} M$ is Totally Non-Cohomologous to Zero, i. e. $H^*(i)$ is onto and that $H^k(\Omega M)$ is a torsion free \mathbb{k} -module for each $k \geq 1$. Then $\chi(M) = 0$ in \mathbb{k} or M is a point.*

Proof. Suppose that $\chi(M)$ is not equal to zero in \mathbb{k} . By part 3) of Theorem 27, $\text{Ker } \sigma^* \cup \chi(M)ev^*(\omega) = \{0\}$. So by Lemma 19, $\tilde{H}^*(\Omega M) = \{0\}$. This means that $H^{>0}(\Omega M) = \{0\}$ and that $\pi_1(M) = \{0\}$. So $H^*(M) \cong \mathbb{k}$. Since $H^{\dim M}(M) = \mathbb{k}\omega$, M must be of dimension 0, i. e. M is a point. \square

Interpretation and proofs of Corollaries 38 and 39 in term of integration along the basis. Let $F \xrightarrow{i} E \xrightarrow{p} M$ be a fibration. Suppose that $\pi_1(M)$ acts trivially on $H^q(F)$. Let $\int i : H^q(F) \rightarrow H^{q+m}(E)$ be the composite

$$H^q(F) \xrightarrow{\omega \otimes Id} H^m(M) \otimes H^q(F) = E_2^{m,q} \twoheadrightarrow E_\infty^{m,q} = F^m H^{m+q}(E) \subset H^{m+q}(E).$$

If M is a sphere, this *integration along the basis* $\int i$ appears in Wang exact sequence and the following two properties are well known [32, Theorems (3.1) and (3.5) Chapter VII]: for $x \in H^*(F)$ and $y \in H^*(E)$,

$$\left(\int i \right) (i^*(y) \cup x) = y \cup \left(\int i \right) (x)$$

and $p^*(\omega) = \left(\int i \right) (1)$. In general, these properties are easy to deduce from the multiplicativity and the naturality of the Serre spectral sequence. In particular, we have $\left(\int i \right) \circ i^*(y) = y \cup p^*(\omega)$. Since the

inclusion of the fibre $i : F \hookrightarrow E$ is the pull-back along p of the embedding $* \hookrightarrow M$, following Section 3, one can define a shriek map $i^!$ for i . In this paper, we will not use that $i^!$ coincides with $(\int i)$ although this should follow from the diagram in [18, (2) p. 12].

Consider the free loop fibration $\Omega M \xrightarrow{i} LM \xrightarrow{ev} M$. In this case, $i_!$ is the intersection morphism $H_{*+\dim M}(LM) \rightarrow H_*(\Omega M)$ defined by Chas and Sullivan.

Let f be an element as in Corollary 38. By Theorem 33,

$$\chi(M) \int i(f) = \chi(M) (\int i) \circ i^*(c) = \chi(M) c \cup ev^*(\omega) = 0.$$

So we have recover Corollary 38.

Suppose that i^* is onto and that $H^*(\Omega M)$ is torsion free, the Serre spectral sequence collapses at the E_2 -term. So $\int i$ is injective. Therefore $\chi(M)f = 0$. And we have recover Corollary 39.

12. TNCZ FREE LOOP FIBRATIONS

Recall our result on TNCZ free loop fibration.

Corollary 40. *(Corollary 39) Let M be a connected, closed \mathbb{k} -oriented manifold. Suppose that the free loop fibration $\Omega M \xrightarrow{i} LM \xrightarrow{ev} M$ is Totally Non-Cohomologous to Zero, i. e. $H^*(i)$ is onto and that $H^k(\Omega M)$ is a torsion free \mathbb{k} -module for each $k \geq 1$. Then $\chi(M) = 0$ in \mathbb{k} or M is a point.*

1) The first examples to have in mind are connected compact Lie groups.

2) Let M be a sphere S^d or the complex or quaternionic projective space $\mathbb{C}\mathbb{P}^n, \mathbb{H}\mathbb{P}^n$. Since $\chi(S^d) = 1 + (-1)^d$ and $\chi(\mathbb{C}\mathbb{P}^n) = \chi(\mathbb{H}\mathbb{P}^n) = n + 1$, it follows from our calculations in [20] that over any commutative ring \mathbb{k} , $H^*(i; \mathbb{k})$ is onto if and only if $\chi(M) = 0$ in \mathbb{k} (when \mathbb{k} is a field, this follows easily from the formality of M using the Jones isomorphism between Hochschild homology and free loop space cohomology).

3) The converse of Corollary 39 is not true: if $n + 1$ is not equal to 0 in \mathbb{k} , take for example $M = \mathbb{C}\mathbb{P}^n \times S^3$.

Over \mathbb{Q} , Vigué-Poirrier has characterised which free loop fibrations are TNCZ.

Theorem 41. [31] *Let X be a simply-connected topological space such that for all n , $H^n(X; \mathbb{Q})$ is finite dimensional. Then*

$$H^*(i; \mathbb{Q}) : H^*(LX; \mathbb{Q}) \rightarrow H^*(\Omega X; \mathbb{Q})$$

is onto if and only if $H^(X; \mathbb{Q})$ is a free graded commutative algebra.*

In [16, Theorem 2], Kuribayashi studied TNCZ free loop fibrations for some homogeneous spaces over a prime field \mathbb{F}_p .

Let $V_k(\mathbb{R}^n)$ denotes the real Stiefel manifold of orthonormal k -frames in \mathbb{R}^n . Using the fibration $S^{n-k} \hookrightarrow V_k(\mathbb{R}^n) \twoheadrightarrow V_{k-1}(\mathbb{R}^n)$, we see that if $k \geq 2$, $\chi(V_k(\mathbb{R}^n)) = 0$. Similarly for the complex or quaternionic Stiefel manifold, $\chi(V_k(\mathbb{C}^n)) = \chi(V_k(\mathbb{H}^n)) = 0$. The Euler characteristic $\chi(G/H)$ of the quotient of a compact connected Lie group G by a connected closed subgroup H of same rank is the quotient $|W(G)|/|W(H)|$ of the cardinals of their Weyl groups [23, VII.Theorem 3.13]. Therefore $\chi(Sp(n)/U(n) = 2^n$ and for Grassmannians $\chi(G_k(\mathbb{C}^n)) = \chi(G_k(\mathbb{H}^n)) = \binom{n}{k}$. We can now rewrite the main theorem of [16] in term of Euler characteristic.

Theorem 42. ([16, Theorem 2])

- 1) Let M be $Sp(n)/U(n)$ or $V_k(\mathbb{C}^n)$ or $V_k(\mathbb{H}^n)$. Then $H^*(i; \mathbb{F}_p) : H^*(LM; \mathbb{F}_p) \rightarrow H^*(\Omega M; \mathbb{F}_p)$ is onto if and only if $\chi(M) = 0$ modulo p .
- 2) Let M be $G_m(\mathbb{C}^{m+n})$ or $G_m(\mathbb{H}^{m+n})$ with m and $n \geq 2$ and p any prime. Then $H^*(i; \mathbb{F}_p) : H^*(LM; \mathbb{F}_p) \rightarrow H^*(\Omega M; \mathbb{F}_p)$ is not onto.
- 3) Let p an odd prime. Then $H^*(i; \mathbb{F}_p) : H^*(LV_m(\mathbb{R}^{m+n}); \mathbb{F}_p) \rightarrow H^*(\Omega V_m(\mathbb{R}^{m+n}); \mathbb{F}_p)$ is onto if and only if n is odd.

So over \mathbb{F}_p , it is not clear when the converse of Corollary 39 holds or not.

Let X be a topological space. Suppose that $\forall n \geq 0$, $H_n(\Omega X; \mathbb{Z})$ is a finitely generated free abelian group, that $H_*(LX; \mathbb{Z})$ has no p -torsion and that $H^*(i; \mathbb{F}_p) : H^*(LX; \mathbb{F}_p) \rightarrow H^*(\Omega X; \mathbb{F}_p)$ is onto. Then by the universal coefficient theorem for homology, $H^*(i; \mathbb{Q}) : H^*(LX; \mathbb{Q}) \rightarrow H^*(\Omega X; \mathbb{Q})$ is onto.

We now give a last result on TNCZ free loop fibration due to Iwase in the context of classifying space BG of finite loop spaces.

Theorem 43. [13, Theorem 2.2] *Let X be a pointed topological space. If $H_*(i; \mathbb{k}) : H_*(\Omega X; \mathbb{k}) \rightarrow H_*(LX; \mathbb{k})$ is injective then the Pontryagin algebra $H_*(\Omega X; \mathbb{k})$ is graded commutative (in particular $\pi_1(X)$ is abelian).*

Proof. We have the following two strictly commutative squares and the following triangle commuting up to a homotopy H .

$$\begin{array}{ccc}
 \Omega X \times \Omega X & \xrightarrow{j} & LX \times_X LX \\
 \mu \downarrow & & \downarrow \mu \\
 \Omega X & \xrightarrow{i} & LX
 \end{array}
 \quad
 \begin{array}{ccc}
 \Omega X \times \Omega X & \xrightarrow{j} & LX \times_X LX \\
 \tau \downarrow & & \downarrow \tau \\
 \Omega X \times \Omega X & \xrightarrow{j} & LX \times_X LX \xrightarrow{\mu} LX
 \end{array}$$

where the maps μ are the composition of loops and the maps τ are the exchange isomorphisms. The homotopy H is the restriction to $[0, 1/2] \times LX \times_X LX$ of the composite of

$$Id \times \mu : S^1 \times LX \times_X LX \rightarrow S^1 \times LX$$

and of the action of the circle on free loops $S^1 \times LX \rightarrow LX$. So finally, $i \circ \mu$ is homotopic to $i \circ \mu \circ \tau$. Since $H_*(i)$ is injective, $H_*(\mu) \circ H_*(\tau) = H_*(\mu)$. \square

Note that our homotopy between $i \circ \mu$ and $i \circ \mu \circ \tau$ is much simpler than the one arriving in $EG \times_G G^{ad}$ given by Iwase in [13, Proof of Lemma 3.1].

Example 44. (Suspension) Suppose that \mathbb{k} is a field. Let X be a path-connected space such that $H_*(X)$ is not concentrated in degree 0. By Bott-Samelson theorem, the Pontryagin algebra $H_*(\Omega\Sigma X)$ is isomorphic to the tensor algebra $TH_+(X)$ on the homology of X in positive degrees. Suppose that $H^*(i; \mathbb{k}) : H^*(L\Sigma X; \mathbb{k}) \rightarrow H^*(\Omega\Sigma X; \mathbb{k})$ is surjective. Then $H_*(i; \mathbb{k}) : H_*(\Omega\Sigma X; \mathbb{k}) \rightarrow H_*(L\Sigma X; \mathbb{k})$ is injective. So by Theorem 43, the Pontryagin ring $H_*(\Omega\Sigma X)$ is graded commutative. So $H_+(X)$ is of dimension 1 and is concentrated in even degree if the characteristic of \mathbb{k} is different from 2 (See [17, Example 2.6] for a proof using Hochschild homology). In particular $\chi(\Sigma X) = 0$ modulo the characteristic of \mathbb{k} .

Conjecture 45. *Let X be a simply-connected finite CW-complex and suppose that \mathbb{k} is a field. If $H^*(i; \mathbb{k}) : H^*(LX; \mathbb{k}) \rightarrow H^*(\Omega X; \mathbb{k})$ is onto then $\chi(X)$ is zero modulo the characteristic of \mathbb{k} or $H_*(X) \cong \mathbb{k}$.*

It follows from the theorem of Vigue-Poirrier recalled above (Theorem 41) that the conjecture is true over the rationals. In Example 44, we have checked the conjecture for suspensions. In this paper, we proved the conjecture when X is a smooth connected, closed \mathbb{k} -oriented manifold M (Corollary 39). We believe that conjecture 45 can be proved easily using Spanier-Whitehead duality.

13. THE RELATIVE FREE LOOPS CASE

Let $g : N \rightarrow M$ be a map. Let g^*LM denote the *relative free loops space* of g which is obtained by the following pull-back

$$\begin{array}{ccc} g^*LM & \longrightarrow & LM \\ p \downarrow & & \downarrow ev \\ N & \xrightarrow{g} & M. \end{array}$$

Theorem 46. *Let $\sigma : N \hookrightarrow g^*LM$, $n \mapsto (n, \text{constant loop } g(n))$ be the section of the projection map $p : g^*LM \rightarrow N$, $(n, w) \mapsto n$. Let M be a smooth h^* -oriented manifold of dimension m . Let $e_{TM} \in h^m(M)$ be the Euler class of the tangent bundle of M . Suppose that g is the composite*

$$N \xrightarrow{g_1} L \xrightarrow{g_2} M$$

where $\begin{cases} a) L \text{ is a smooth manifold without boundary and } g_2 \text{ is smooth and} \\ b) g_1 \text{ is a (Serre) fibration.} \end{cases}$

Then for any $b \in h^*(g^*LM)$,

$$b \cup p^* \circ g^*(e_{TM}) = p^* \circ \sigma^*(b) \cup p^* \circ g^*(e_{TM}).$$

In particular if there is an integer $\dim N$ such that $\forall i > \dim N$, $h^i(N) = \{0\}$ then for all $b \in h^*(g^*LM)$ of degree $|b| > \dim N - m$, $b \cup p^* \circ g^*(e_{TM}) = 0$.

Example 47. An interesting example is when the fibration $g = g_1 : \text{map}(S, M) \rightarrow M$ is the evaluation at the base point of a well-pointed space S . In this case, the pull-back g^*LM of LM along g is the space $\text{map}(S \vee S^1, M)$ of maps from the wedge of S and the circle to M .

Example 48. A generalization of the preceding example is when $g_2 : L \rightarrow M$ is any smooth map and the fibration $g_1 : \text{map}(S, L) \rightarrow L$ is the evaluation at the base point of a well-pointed space S . In this case, the pull-back g^*LM of LM along g is the space $\text{map}((S \vee S^1, S), (M, L))$

of couples of maps (φ, ψ) such that the square

$$\begin{array}{ccc} S & \xrightarrow{\psi} & L & \text{com-} \\ \downarrow & & \downarrow g_2 & \\ S \vee S^1 & \xrightarrow{\varphi} & M. & \end{array}$$

mutates.

Corollary 49. *Let M be a smooth h^* -oriented manifold of dimension m . Let $e_{TM} \in h^m(M)$ be the Euler class of the tangent bundle of M . Denote by $\vee_n S^1$ the wedge of $n \geq 0$ circles. Let $\sigma_n : M \hookrightarrow \text{map}(\vee_n S^1, M)$, $m \mapsto \text{constant map } m$ be the section of the evaluation map $ev_n : \text{map}(\vee_n S^1, M) \rightarrow M$. Then for any $b \in h^*(\text{map}(\vee_n S^1, M))$,*

$$b \cup ev_n^*(e_{TM}) = ev_n^* \circ \sigma_n^*(b) \cup ev_n^*(e_{TM}).$$

Proof. When $n = 0$, the formula is true since ev_0 and σ_0 are just the identity map of M . By induction on n , suppose that for any $a \in h^*(\text{map}(\vee_{n-1} S^1, M))$,

$$a \cup ev_{n-1}^*(e_{TM}) = ev_{n-1}^* \circ \sigma_{n-1}^*(a) \cup ev_{n-1}^*(e_{TM}).$$

By example 47 in the case $S = \vee_{n-1} S^1$ and $g = ev_{n-1}$, for any $b \in h^*(\text{map}(S \vee S^1, M))$,

$$b \cup p^* \circ ev_{n-1}^*(e_{TM}) = p^* (\sigma^*(b) \cup ev_{n-1}^*(e_{TM})).$$

By taking $a = \sigma^*(b)$, the latter is equal to

$$p^* (ev_{n-1}^* \circ \sigma_{n-1}^* \circ \sigma^*(b) \cup ev_{n-1}^*(e_{TM}))$$

Since $p^* \circ ev_{n-1}^* = ev_n^*$ and $\sigma_{n-1}^* \circ \sigma^* = \sigma_n^*$, the conclusion follows. \square

Remark 50. Again, let S be a well-pointed space. Suppose that the fibration $\text{map}_*(S \vee S^1, M) \xrightarrow{i} \text{map}(S \vee S^1, M) \xrightarrow{ev} M$ is Totally Non-Cohomologous to Zero, i. e. $H^*(i)$ is onto. Let $\pi : S \vee S^1 \rightarrow S^1$ be the canonical projection. Then we have the commutative triangle

$$\begin{array}{ccc} LM & \xrightarrow{\text{map}(\pi, M)} & \text{map}(S \vee S^1, M) \\ & \searrow ev & \downarrow ev \\ & & M \end{array}$$

Since the induced map between the fibers in cohomology, $H^*(\text{map}_*(\pi, M)) : H^*(\text{map}_*(S \vee S^1, M)) \rightarrow H^*(\Omega M)$ is surjective, the free loop fibration $\Omega M \xrightarrow{i} LM \xrightarrow{ev} M$ is also Totally Non-Cohomologous to Zero, i. e. $H^*(i)$ is onto.

Conclusion: the preceding corollary together with Lemma 19 is not really interesting to see if the fibration $(\Omega M)^{\times n} \xrightarrow{i} \text{map}(\vee_n S^1, M) \xrightarrow{ev_n} M$ is Totally Non-Cohomologous to Zero or not.

Let $f : N \rightarrow M$ and $g : N \rightarrow M$ be two maps. Let $N \times_f M^I \times_g N$ denote the *homotopy coincidence point space* of f and g which is obtained by the following pull-back

$$\begin{array}{ccc} N \times_f M^I \times_g N & \longrightarrow & M^I \\ p \downarrow & & \downarrow (ev_0, ev_1) \\ N & \xrightarrow{(f, g)} & M \times M. \end{array}$$

Lemma 51. Let $\bar{\xi} : N \times_f M^I \times_g N \hookrightarrow {}^{1, f}(N \times M)^{1, g}$ be the map from the homotopy coincidence point space of f and g to the homotopy fibre product of $(1, f) : N \rightarrow N \times M$ and $(1, g) : N \rightarrow N \times M$ defined by

$$\bar{\xi}(n, \omega) = (n, \text{the path } t \mapsto (n, \omega(t)), n)$$

for any $n \in N$ and any path $\omega : I \rightarrow M$ such that $\omega(0) = f(n)$ and $\omega(1) = g(n)$. Then $\bar{\xi}$ is a homotopy equivalence

Proof. Consider the three pull-back squares

$$\begin{array}{ccccc}
 {}^{1,f}(N \times M)^{1,g} & \longrightarrow & (N \times M)^I \times_{1,g} N & \longrightarrow & (N \times M)^I \\
 (ev_0, ev_1) \downarrow & & (ev_0, ev_1) \downarrow & & \downarrow (ev_0, ev_1) \\
 N \times N & \xrightarrow{(1,f) \times 1} & (N \times M) \times N & \xrightarrow{1 \times (1,g)} & (N \times M) \times (N \times M) \\
 p_1 \downarrow & & \downarrow p_1 & & \\
 N & \xrightarrow{(1,f)} & N \times M & &
 \end{array}$$

Here p_1 are the projections on the first factor. Consider also the two pull-back squares

$$\begin{array}{ccccc}
 N \times_f M^I \times_g N & \longrightarrow & M^I \times_g N & \longrightarrow & M^I \\
 p \downarrow & & (p_N, ev_0) \downarrow & & (ev_1, ev_0) \downarrow \\
 N & \xrightarrow{(1,f)} & N \times M & \xrightarrow{g \times 1} & M \times M
 \end{array}$$

Let $\xi : M^I \times_g N \hookrightarrow (N \times M)^I \times_{1,g} N$ be the map defined by

$$\xi(n, \omega) = (\text{the path } t \mapsto (n, \omega(t)), n)$$

for any $n \in N$ and any path $\omega : I \rightarrow M$ such that $\omega(1) = g(n)$. Obviously ξ is a homotopy equivalence. We obtain the following commutative diagram where the two squares are pull-backs according to the two previous diagrams.

$$\begin{array}{ccc}
 N \times_f M^I \times_g N & \longrightarrow & M^I \times_g N \\
 \downarrow \bar{\xi} & & \approx \downarrow \xi \\
 {}^{1,f}(N \times M)^{1,g} & \longrightarrow & (N \times M)^I \times_{1,g} N \\
 \downarrow ev_0 & & \downarrow ev_0 \\
 N & \xrightarrow{(1,f)} & N \times M
 \end{array}$$

p (left curved arrow), (p_N, ev_0) (right curved arrow)

By decomposing $(1, f)$ into the composite of a homotopy equivalence and of a fibration, we show using the structure of model category on topological spaces, that $\bar{\xi}$ is a homotopy equivalence since both ev_0 and $(p_N, ev_0) = ev_0 \circ \xi$ are fibrations. \square

Proof of Theorem 46. Consider the two pull-back squares

$$\begin{array}{ccc}
 N & \xrightarrow{(1,g)} & N \times M \\
 g_1 \downarrow & & \downarrow g_1 \times 1 \\
 L & \xrightarrow{(1,g_2)} & L \times M \\
 g_2 \downarrow & & \downarrow g_2 \times 1 \\
 M & \xrightarrow{\Delta} & M \times M.
 \end{array}$$

Since g_2 and $1 : M \rightarrow M$ are transverse, $g_2 \times 1$ is transverse to the diagonal embedding Δ . And so by Remark 9, $(1, g_2) : L \rightarrow L \times M$ is a proper embedding of codimension m with h^* -oriented normal bundle.

Since g_1 is a (Serre) fibration, $g_1 \times 1$ is also a (Serre) fibration. Therefore $(1, g) : N \rightarrow N \times M$ is the pull-back of an embedding in the sense of definition 8. So we can apply part 4) of Theorem 12. Since the homotopy equivalence $\bar{\xi} : N \times_f M^I \times_g N \xrightarrow{\cong} {}^1, f(N \times M)^{1, g}$ of Lemma 51 in the case $f = g$ commutes with the projection maps, i. e. $ev_0 \circ \bar{\xi} = p$ and also with the two sections σ , the ideal $\text{Ker } \sigma^* : h^*(g^*LM) \rightarrow h^*(N)$ satisfies

$$\text{Ker } \sigma^* \cup p^* \circ g^*(e_{TM}) = \{0\}.$$

□

REFERENCES

- [1] Marcelo Aguilar, Samuel Gitler, and Carlos Prieto, *Algebraic topology from a homotopical viewpoint*, Universitext, Springer-Verlag, New York, 2002.
- [2] K. S. Brown, *Cohomology of groups*, Graduate Texts in Mathematics, no. 87, Springer-Verlag, New York, 1994.
- [3] M. Chas and D. Sullivan, *String topology*, preprint: math.GT/9911159, 1999.
- [4] David Chataur and Jean-Claude Thomas, *Frobenius rational loop algebra*, Manuscripta Math. **122** (2007), no. 3, 305–319.
- [5] Ralph L. Cohen and Véronique Godin, *A polarized view of string topology*, Topology, geometry and quantum field theory, London Math. Soc. Lecture Note Ser., vol. 308, Cambridge Univ. Press, Cambridge, 2004, pp. 127–154.
- [6] Ralph L. Cohen, Kathryn Hess, and Alexander A. Voronov, *String topology and cyclic homology*, Birkhäuser Verlag, Basel, 2006.
- [7] Eldon Dyer, *Cohomology theories*, Mathematics Lecture Note Series, W. A. Benjamin, Inc., New York-Amsterdam, 1969.
- [8] Yves Félix and Jean-Claude Thomas, *String topology on Gorenstein spaces*, Math. Ann. **345** (2009), no. 2, 417–452.
- [9] Mark Goresky and Nancy Hingston, *Loop products and closed geodesics*, Duke Math. J. **150** (2009), no. 1, 117–209.
- [10] A. Hatcher, *Algebraic topology*, Cambridge University Press, 2002.
- [11] ———, *Spectral Sequence in Algebraic Topology*, <http://www.math.cornell.edu/hatcher/SSAT/SSATpage.html>, 2004.

STRING TOPOLOGY, EULER CLASS AND TNCZ FREE LOOP FIBRATIONS

- [12] Morris W. Hirsch, *Differential topology*, Springer-Verlag, New York, 1976, Graduate Texts in Mathematics, No. 33.
- [13] Norio Iwase, *Adjoint action of a finite loop space*, Proc. Amer. Math. Soc. **125** (1997), no. 9, 2753–2757.
- [14] Sadok Kallel and Paolo Salvatore, *Rational maps and string topology*, Geom. Topol. **10** (2006), 1579–1606.
- [15] Katsuo Kawakubo, *The theory of transformation groups*, japanese ed., The Clarendon Press Oxford University Press, New York, 1991.
- [16] Katsuhiko Kuribayashi, *On the mod p cohomology of the spaces of free loops on the Grassmann and Stiefel manifolds*, J. Math. Soc. Japan **43** (1991), no. 2, 331–346.
- [17] ———, *Module derivations and non triviality of an evaluation fibration*, Homology Homotopy Appl. **4** (2002), no. 1, 87–101.
- [18] J-F Le Borgne, *String spectral sequence*, preprint: math.AT/0409597, 2004.
- [19] John M. Lee, *Introduction to smooth manifolds*, second ed., Graduate Texts in Mathematics, vol. 218, Springer, New York, 2013.
- [20] L. Menichi, *The cohomology ring of free loop spaces*, Homology Homotopy Appl. **3** (2001), no. 1, 193–224.
- [21] ———, *On the cohomology algebra of a fiber*, Algebr. Geom. Topol. **1** (2001), 719–742.
- [22] John W. Milnor and James D. Stasheff, *Characteristic classes*, Princeton University Press, Princeton, N. J., 1974, Annals of Mathematics Studies, No. 76.
- [23] Mamoru Mimura and Hirosi Toda, *Topology of Lie groups. I, II*, vol. 91, American Mathematical Society, Providence, RI, 1991.
- [24] Edwin H. Spanier, *Algebraic topology*, Springer-Verlag, New York, 1981.
- [25] A. Stacey, *The differential topology of loop spaces*, preprint: math.DG/0510097, 2005.
- [26] Dennis Sullivan, *Open and closed string field theory interpreted in classical algebraic topology*, Topology, geometry and quantum field theory, London Math. Soc. Lecture Note Ser., vol. 308, Cambridge Univ. Press, Cambridge, 2004, pp. 344–357.
- [27] H. Tamanoi, *TQFT string operations in open-closed string topology*, ArXiv e-prints (2008).
- [28] Hirotaka Tamanoi, *Cap products in string topology*, Algebr. Geom. Topol. **9** (2009), no. 2, 1201–1224.
- [29] ———, *Loop coproducts in string topology and triviality of higher genus TQFT operations*, J. Pure Appl. Algebra **214** (2010), no. 5, 605–615.
- [30] Tammo tom Dieck, *Algebraic topology*, EMS Textbooks in Mathematics, European Mathematical Society (EMS), Zürich, 2008.
- [31] Micheline Vigué-Poirrier, *Dans le fibré de l'espace des lacets libres, la fibre n'est pas, en général, totalement non cohomologue à zéro*, Math. Z. **181** (1982), no. 4, 537–542.
- [32] G. Whitehead, *Elements of homotopy theory*, Graduate Texts in Mathematics, vol. 61, Springer-Verlag, 1978.

LAREMA, UMR CNRS 6093, UNIVERSITÉ D'ANGERS, 2 BD LAVOISIER, 49045
ANGERS, FRANCE

E-mail address: `firstname.lastname at univ-angers.fr`